

VERIFICATION OF TRANSLATION

I, Norio Ohmori, registered Patent Attorney, having my business place at c/o Abe, Ikubo and Katayama, Fukuoka Building, 9th Floor, 8-7, Yaesu 2-Chome, Chuo-ku, Tokyo 104-0028 Japan, state the following:


I am fluent in both the English and Japanese languages and capable of translating documents from one into the other of these languages.

The attached document is a true and accurate English translation to the best of my knowledge and belief of:

Japanese Patent Application No. 2004-48593

I state that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true.

Signature:

A handwritten signature in black ink, appearing to read 'Norio Ohmori', written over a horizontal line.

Norio Ohmori

Date : November 19, 2009

[Title of Document] CLAIMS

[Claim 1]

A gene consisting of at least one of the following definitions correlated with prediction of the postoperative prognosis of breast cancer;

5 1) a marker gene group capable of establishing classification of genes from breast cancer patients died within 5 years after a surgical operation (5y-D group) and genes from patients survived free of disease for several years or more after the operation (5y-S group), depending on their expression functions, in estrogen receptor-negative breast cancer,

2) a marker gene group capable of establishing classification of genes from n0 breast
10 cancer patients recurred within 5 years after an operation (5Y-R group) and genes from patients survived free of disease for 5 years or more after the operation (5Y-F group), depending on their expression functions, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in the operation,

3) a marker gene group capable of establishing classification of genes from breast
15 cancer patients died within 5 years after a surgical operation (5D group) and genes from patients survived free of disease for several years or more after the operation (5S group), depending on their expression functions, in primary breast cancer.

[Claim 2]

A gene selected from the following sequences correlated with prediction of the
20 postoperative prognosis of primary breast cancer;

pro-alpha-1 type 3 collagen (PIIIP),

complement component C1r,

dihydropyrimidinase-like 3 (DPYSL3),

protein tyrosine kinase 9-like (PTK9L),

25 carboxypeptidase E (CPE),

alpha-tubulin,

beta-tubulin,

heat shock protein HSP 90-alpha gene,

malate dehydrogenase,

NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (NDUFB3).

[Claim 3]

5 A gene selected from the following sequences highly expressed in a group of good
prognosis correlated with prediction of the postoperative prognosis of primary breast cancer;
pro-alpha-1 type 3 collagen (PIIIP),
complement component C1r,
dihydropyrimidinase-like 3 (DPYSL3),
protein tyrosine kinase 9-like (PTK9L),
10 carboxypeptidase E (CPE),
alpha-tubulin,
beta-tubulin.

[Claim 4]

15 A gene selected from the following sequences highly expressed in a group of bad
prognosis correlated with prediction of the postoperative prognosis of primary breast cancer;
heat shock protein HSP 90-alpha gene,
malate dehydrogenase,
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (NDUFB3).

[Claim 5]

20 A gene selected from the following sequences correlated with prediction of the
postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph
node in operation;
AF058701/ DNA polymerase zeta catalytic subunit (REV3),
AI066764/ lectin, galactoside-binding, soluble, 1 (galectin 1),
25 x15940/ ribosomal protein L31.,
Hs.94653/ neurochondrin (KIAA0607),
M13436/ ovarian beta-A-inhibin,
Hs.5002/ copper chaperone for superoxide dismutase; CCS,

D67025/ proteasome (prosome, macropain) 26S subunit, non-ATPase, 3,
M80469/ MHC class I HLA-J gene,
Hs.4864/ ESTs,
Hs.106326/ ESTs.

5 [Claim 6]

A gene selected from the following sequences highly expressed in a group of bad prognosis correlated with prediction of the postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in operation;

AF058701/ DNA polymerase zeta catalytic subunit (REV3),

- 10 AI066764/ lectin, galactoside-binding, soluble, 1 (galectin 1),
x15940/ ribosomal protein L31.

[Claim 7]

A gene selected from the following sequences highly expressed in a group of good prognosis correlated with prediction of the postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in operation;

Hs.94653/ neurochondrin (KIAA0607),

M13436/ ovarian beta-A-inhibin,

Hs.5002/ copper chaperone for superoxide dismutase; CCS,

D67025/ proteasome (prosome, macropain) 26S subunit, non-ATPase, 3,

- 20 M80469/ MHC class I HLA-J gene,

Hs.4864/ ESTs,

Hs.106326/ ESTs.

[Claim 8]

- 25 A gene selected from the following sequences correlated with prediction of the postoperative prognosis, in estrogen receptor-negative breast cancer;

Hs.108504/ FLJ20113/ ubiquitin-specific protease otubain 1

Hs.146550/ MYH9/ myosin, heavy polypeptide 9, non-muscle

Hs.194691/ RAI3/ retinoic acid induced 3

Hs.1975/ TDRD3/ tudor domain containing 3

Hs.203952/ TRRAP/ transformation/transcription domain-associated protein

Hs.278607/ GSA7/ ubiquitin activating enzyme E1-like protein

Hs.429/ ATP5G3/

- 5 ATP synthase, H⁺ transporting, mitochondrial F₀ complex, subunit c (subunit 9) isoform 3

Hs.75305/ AIP/ aryl hydrocarbon receptor interacting protein

Hs.81170/ PIM1/ pim-1 oncogene

Hs.99987/ ERCC2/

- 10 excision repair cross-complementing rodent repair deficiency, complementation group 2

Y12781/ Transducin (beta) like 1 protein

Hs.104417/ KIAA1205 protein

cl.21783/ Hypothetical protein

- 15 Hs.112628/ Hypothetical protein: MGC43581

Hs.170345/ Hypothetical protein FLJ13710

Hs.53996/ weakly similar to zinc finger protein 135

Hs.55422/ Hypothetical protein

Hs.112718/ EST

- 20 Hs.115880/ EST

Hs.126495/ EST.

[Claim 9]

A gene selected from Claim 8, as a gene highly expressed in a group of bad prognosis.

- 25 [Claim 10]

A DNA microarray carrying thereon the gene according to any one of Claims 1 to 9.

[Claim 11]

The microarray according to Claim 10, wherein the DNA microarray is a fiber type

microarray.

[Claim 12]

A method of inspecting the postoperative prognosis of breast cancer using as a marker the gene and/or probe according to any one of Claims 1 to 9.

5 [Claim 13]

A method of inspecting the postoperative prognosis of breast cancer using the microarray according to Claim 10 or 11.

[Claim 14]

10 A method of screening cancer therapeutic medicines for controlling the postoperative prognosis of breast cancer using as a marker the gene and/or probe according to any one of Claims 1 to 9.

[Claim 15]

A method of screening cancer therapeutic medicines for controlling the postoperative prognosis of breast cancer using the microarray according to Claim 10 or 11.

15 [Claim 16]

A diagnosis kit for the postoperative prognosis of breast cancer containing a reagent using as a marker the gene and/or probe according to any one of Claims 1 to 9.

[Claim 17]

The diagnosis kit according to Claim 16, wherein the kit comprises a microarray.

20 [Claim 18]

The diagnosis kit according to Claim 17, wherein the microarray is a fiber type microarray.

[Title of Document] SPECIFICATION

[Title of the Invention] GENE RELATING TO ESTIMATION OF POSTOPERATIVE
PROGNOSIS FOR BREAST CANCER

[TECHNICAL FIELD]

5 [0001]

The present invention relates to a gene correlated with prediction of the postoperative prognosis of breast cancer. Further, the present invention relates to a method of inspecting the postoperative prognosis of breast cancer using this gene, a method of screening cancer therapeutic medicines for controlling the postoperative prognosis of breast cancer, and a
10 diagnosis kit for the postoperative prognosis of breast cancer.

[BACKGROUND ART]

[0002]

Breast cancer is a disease situated as a superior cause for female lethality due to cancer, however, there are found still no dominant reasons for determining the grade of malignancy
15 and survival prognosis from the biological standpoint.

[0003]

The condition of an estrogen receptor (ER) is one determining element for clinical and biological symptoms of human breast cancer. Adjuvant hormone therapeutics is usually effective in ER-positive breast cancer patients irrespective of age, condition in the
20 menopause, correlation with axillary nodes, and tumor diameter. However, ER-negative breast cancer is resistance to this therapeutic method (Non Patent Literatures No. 1 and 2). Patients having an ER-negative tumor do not necessarily show the same response to chemical therapy. Since existent indices cannot classify breast cancer of this type depending on clinical symptom, the postoperative prognosis is recognized to be various
25 (Non Patent Literatures No. 3 and 4).

[0004]

Prognosis of breast cancer patients with no lymph node metastasis (node-negative breast cancer; n0) is better than that of metastatic breast cancer patients. However, in

Japan, the present inventors have found that 16% of node-negative breast cancer patients relapse within 5 years after the initial operation (Non Patent Literature No. 5).

[0005]

Prediction of the postoperative prognosis of breast cancer patients shows increasing in
5 importance from the standpoint of adjuvant therapeutics currently utilizable. A gene
marker which is useful in identifying patients showing a possibility of relapsing after an
operation gives a merit which suitable preoperative adjuvant therapeutics can be applied to a
high risk patient, and enables prevention of occurrence of unnecessary, complicated and
uncomfortable side effects.

10 [0006]

Conventionally, postoperative procedures for individual patients are determined
depending on tumor diameter and the stage, metastasis to a lymph node, diagnosis by
clinicopathological factors, search of a hormone receptor, and the like, however, they are not
critical methods (Non Patent Literatures No. 6-11).

15 [0007]

Recently, there is a prognosis marker for postoperative breast cancer patients,
intending determination of an importance of mutations of genes. These gene mutations
include a mutation of p53 (Non Patent Literature No. 12), loss of heterozygosity in several
alleles (Non Patent Literature No. 13), and abnormal expressions of a BRCA2 gene (Non
20 Patent Literature No. 14), WT1 gene (Non Patent Literature No. 15), HER2/neu gene (Non
Patent Literature No. 16) and Ki-67 gene (Non Patent Literature No. 17). However, these
would not be recognized as effective prognosis predicting means when taking into
consideration a fact which a cancer is a disease owing to accumulation of abnormalities of
multiple genes.

25 [0008]

Further, in these years, genome projects in various organisms are being progressed,
and a lot of genes and their base sequences typically including a human gene are being
clarified quickly. The function of a gene having a clarified sequence can be checked by

various methods. As one of the effective methods, known is a gene expression analysis method utilizing clarified base sequence information. For example, there are developed methods utilizing various nucleic acid-nucleic acid hybridization reactions and various PCR reactions as typified by Northern Hybridization, and relations between various genes and expressions of their organism functions can be checked by these methods. Though the number of applicable genes is limited in these methods, there have been developed a methodology and a novel analysis method called DNA microarray method (DNA chip method) enabling lump expression analysis of multiple genes, for carrying out comprehensive and systemic analysis of extremely many genes such as one individual level, as being clarified recently through genome projects.

[0009]

As the DNA microarray, a lot of shapes are known such as that in which DNA synthesis is conducted on many discrete cells applying a lithography technology (Patent Literature No. 1), that in which cells composed of grooves or holes are formed on a board and a probe is fixed to the inner wall of the cell (Patent Literatures No. 2 and 3), a microarray in which a probe is fixed to a gel such as acrylamide and the like for increasing the amount of a probe to be fixed on a chip (Patent Literatures No. 4 and 5), and the like.

[0010]

Some of the present inventors previously developed a new type of a microarray and a method for producing the same (Patent Literatures No. 6-7).

The above invention provides a chip that can be obtained by fabricating a nucleic acid fixed gel retaining fiber array which retains a nucleic acid fixed gel, and cutting this array along a direction crossing the fiber axis of the array. The chip can be utilized as a two-dimensional high-density nucleic acid fixed array, namely, a microarray.

[0011]

Recent studies have found that a cDNA microarray technology is effective for identification of a novel gene marker for cancer diagnosis. To date, some researchers have carried out microarray analysis of breast cancer, however, there is no description about data

of breast cancer gene expression property capable of predicting the postoperative prognosis of breast cancer (Non Patent Literatures No. 18-24). As one exception, it is shown that a specific profile of a lymph node metastasis negative tumor gives a prediction of a short interval before progressing to distant metastasis (Non Patent Literature No. 25).

- 5 [Patent Literature No. 1] USP 5445934
[Patent Literature No. 2] Tokkyo KOKAI (unexamined Japanese patent application) No. 11-108928
[Patent Literature No. 3] Tokkyo KOKAI No. 2000-78998
[Patent Literatures No. 4] USP 5770721
- 10 [Patent Literatures No. 5] Tokkyo KOKAI No. 2000-60554
[Patent Literature No. 6] Tokkyo KOKAI No. 2000-270878
[Patent Literature No. 7] Tokkyo KOKAI No. 2000-270879
[Non Patent Literature No. 1] J Clin Oncology (2001) 19, 3817-1827
[Non Patent Literature No. 2] Breast Cancer (2001) 8, 298-304
- 15 [Non Patent Literature No. 3] J Natl Inst (1991) 83, 154-155
[Non Patent Literature No. 4] J Natl Cancer Inst (2000) 93, 979-989
[Non Patent Literature No. 5] Clin Cancer Res (2000) 6, 3193-3198
[Non Patent Literature No. 6] Cancer (1982) 50, 2131-2138
[Non Patent Literature No. 7] Histopathology (1991) 19, 403-410
- 20 [Non Patent Literature No. 8] Int J Cancer (1996) 69, 135-141
[Non Patent Literature No. 9] Am J Clin Oncol (1997) 20, 546-551
[Non Patent Literature No. 10] Eur J Cancer (2002) 38, 1329-1334
[Non Patent Literature No. 11] Jpn J Cancer Res (2000) 91, 293-300
[Non Patent Literature No. 12] Breast Cancer Res Treat (2001) 69, 65-68), loss of
25 heterozygosity in several alleles
[Non Patent Literature No. 13] Int J Clin Oncol (2001) 6, 6-12
[Non Patent Literature No. 14] Int J Cancer (2002) 198, 879-882
[Non Patent Literature No. 15] Clin Cancer Res (2002) 8, 1167-1171

- [Non Patent Literature No. 16] Arch Surg (2000) 135, 1469-1474
- [Non Patent Literature No. 17] J Pathol (1999) 187, 207-216
- [Non Patent Literature No. 18] Proc Natl Acad Sci USA (1999) 96, 9212-9217
- [Non Patent Literature No. 19] Nature (2000) 406, 747-752
- 5 [Non Patent Literature No. 20] Proc Natl Acad Sci USA (2001) 98, 11462-11467
- [Non Patent Literature No. 21] Cancer Res (2001) 61, 5979-5984
- [Non Patent Literature No. 22] Cancer Res (2000) 60, 2232-2238
- [Non Patent Literature No. 23] Cancer Res (2001) 61, 5168-5178
- [Non Patent Literature No. 24] Proc Natl Acad Sci USA (2001) 98, 10869-10874
- 10 [Non Patent Literature No. 25] N Engl J Med (2002) 347, 1999-2009

[DISCLOSURE OF THE INVENTION]

[Problem to Be Solved]

[0012]

15 The present invention has an object of providing innovative means for predicting the postoperative prognosis of breast cancer patients from the standpoint of gene expression, based on results obtained by genome-wide and comprehensive analysis on gene expression in breast cancer.

[Means to Solve the Problem]

[0013]

20 The present inventors have comprehensively analyzed gene expression of a human gene by a DNA microarray and compared gene expression functions of breast cancers in various conditions, thereby, establishing a system for predicting the postoperative prognosis of breast cancer.

[0014]

25 That is, the present invention provides the following genes (groups) (1) to (8).

(1) A gene consisting of at least one of the following definitions correlated with prediction of the postoperative prognosis of breast cancer;

1) a marker gene group capable of establishing classification of genes from breast

cancer patients died within 5 years after a surgical operation (5y-D group) and genes from patients survived free of disease for several years or more after the operation (5y-S group), depending on their expression functions, in estrogen receptor-negative breast cancer,

2) a marker gene group capable of establishing classification of genes from n0 breast cancer patients recurred within 5 years after an operation (5Y-R group) and genes from patients survived free of disease for 5 years or more after the operation (5Y-F group), depending on their expression functions, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in the operation,

3) a marker gene group capable of establishing classification of genes from breast cancer patients died within 5 years after a surgical operation (5D group) and genes from patients survived free of disease for several years or more after the operation (5S group), depending on their expression functions, in primary breast cancer.

(2) A gene selected from the following sequences correlated with prediction of the postoperative prognosis of primary breast cancer;

15 pro-alpha-1 type 3 collagen (PIIIP),
complement component C1r,
dihydropyrimidinase-like 3 (DPYSL3),
protein tyrosine kinase 9-like (PTK9L),
carboxypeptidase E (CPE),
20 alpha-tubulin,
beta-tubulin,
heat shock protein HSP 90-alpha gene,
malate dehydrogenase,
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (NDUFB3).

25 (3) A gene selected from the following sequences highly expressed in a group of good prognosis correlated with prediction of the postoperative prognosis of primary breast cancer;
pro-alpha-1 type 3 collagen (PIIIP),
complement component C1r,

dihydropyrimidinase-like 3 (DPYSL3),
protein tyrosine kinase 9-like (PTK9L),
carboxypeptidase E (CPE),
alpha-tubulin,
5 beta-tubulin.

(4) A gene selected from the following sequences highly expressed in a group of bad prognosis correlated with prediction of the postoperative prognosis of primary breast cancer; heat shock protein HSP 90-alpha gene,
malate dehydrogenase,

10 NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (NDUFB3).

(5) A gene selected from the following sequences correlated with prediction of the postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in operation;

AF058701/ DNA polymerase zeta catalytic subunit (REV3),
15 AI066764/ lectin, galactoside-binding, soluble, 1 (galectin 1),
x15940/ ribosomal protein L31.,
Hs.94653/ neurochondrin (KIAA0607),
M13436/ ovarian beta-A-inhibin,
Hs.5002/ copper chaperone for superoxide dismutase; CCS,
20 D67025/ proteasome (prosome, macropain) 26S subunit, non-ATPase, 3,
M80469/ MHC class I HLA-J gene,
Hs.4864/ ESTs,
Hs.106326/ ESTs.

(6) A gene selected from the following sequences highly expressed in a group of bad
25 prognosis correlated with prediction of the postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in operation;
AF058701/ DNA polymerase zeta catalytic subunit (REV3),
AI066764/ lectin, galactoside-binding, soluble, 1 (galectin 1),

x15940/ ribosomal protein L31.

(7) A gene selected from the following sequences highly expressed in a group of good prognosis correlated with prediction of the postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in operation;

- 5 Hs.94653/ neurochondrin (KIAA0607),
M13436/ ovarian beta-A-inhibin,
Hs.5002/ copper chaperone for superoxide dismutase; CCS,
D67025/ proteasome (prosome, macropain) 26S subunit, non-ATPase, 3,
M80469/ MHC class I HLA-J gene,
- 10 Hs.4864/ ESTs,
Hs.106326/ ESTs.

(8) A gene selected from the following sequences correlated with prediction of the postoperative prognosis, in estrogen receptor-negative breast cancer;

- Hs.108504/ FLJ20113/ ubiquitin-specific protease otubain 1
- 15 Hs.146550/ MYH9/ myosin, heavy polypeptide 9, non-muscle
Hs.194691/ RAI3/ retinoic acid induced 3
Hs.1975/ TDRD3/ tudor domain containing 3
Hs.203952/ TRRAP/ transformation/transcription domain-associated protein
Hs.278607/ GSA7/ ubiquitin activating enzyme E1-like protein
- 20 Hs.429/ ATP5G3/

ATP synthase, H⁺ transporting, mitochondrial F₀ complex, subunit c (subunit 9) isoform 3

Hs.75305/ AIP/ aryl hydrocarbon receptor interacting protein
Hs.81170/ PIM1/ pim-1 oncogene

- 25 Hs.99987/ ERCC2/
excision repair cross-complementing rodent repair deficiency,
complementation group 2
Y12781/ Transducin (beta) like 1 protein

Hs.104417/ KIAA1205 protein
cl.21783/ Hypothetical protein
Hs.112628/ Hypothetical protein: MGC43581
Hs.170345/ Hypothetical protein FLJ13710

5 Hs.53996/ weakly similar to zinc finger protein 135

Hs.55422/ Hypothetical protein

Hs.112718/ EST

Hs.115880/ EST

Hs.126495/ EST

10 [0015]

The present invention also provides a gene selected from the above-mentioned (8), as a gene highly expressed in a group of bad prognosis.

Further, the present invention provides a DNA microarray carrying thereon the gene according to any one of the above-mentioned (1) to (9), and preferably, the DNA microarray
15 is a fiber type microarray.

[0016]

The above-mentioned gene can be used as a marker gene for postoperative prognosis of breast cancer. Further, it can be also used as a marker gene for cancer therapeutic medicines for controlling the postoperative prognosis of breast cancer.

20 [0017]

The marker gene can be included as a reagent, and can be used as a diagnosis kit. The reagent kit includes a DNA microarray carrying thereon a marker gene, preferably, a fiber type microarray.

[Effect of the Invention]

25 [0018]

According to the means of the present invention, completely novel breast cancer correlated genes have been found and simultaneously, it has been found that these genes are correlated deeply with malignant degeneration of breast cancer and finally, exert an

influence on the prognosis of breast cancer patients. Further, by establishing a mathematical formula for evaluating expression condition of the found gene, a completely novel and effective breast cancer postoperative prognosis predicting system has been developed. The system of the present invention from the standpoint of gene expression is
5 believed to be an innovative prognosis predicting system arresting the biological essentiality of a cancer, utterly different from conventional prognosis evaluation methods, when taking into consideration a fact which a cancer is a disease owing to abnormality of a gene.

[BEST MODE FOR CARRYING OUT THE INVENTION]

10 [0019]

The marker gene group correlated with prediction of the postoperative prognosis of breast cancer as one aspect of the present invention is obtained by analysis by cDNA microarray of the expression functions of genes from patients manifesting death or recurring within 5 years after a surgical operation and patients survived for 5 years or more after the
15 operation, in estrogen receptor-negative breast cancer, node-negative breast cancer and primary breast cancer.

[0020]

Specifically, one aspect of the present invention is a gene consisting of at least one of the following definitions selected from known sequences correlated with prediction of the
20 postoperative prognosis of breast cancer;

- 1) a marker gene group capable of establishing classification of genes from breast cancer patients died within 5 years after a surgical operation (5y-D group) and genes from patients survived free of disease for several years or more after the operation (5y-S group), depending on their expression functions, in estrogen receptor-negative breast cancer,
- 25 2) a marker gene group capable of establishing classification of genes from n0 breast cancer patients recurred within 5 years after an operation (5Y-R group) and genes from patients survived free of disease for 5 years or more after the operation (5Y-F group), depending on their expression functions, in (node-negative)(n0) breast cancer with no

metastasis to a lymph node in the operation,

3) a marker gene group capable of establishing classification of genes from breast cancer patients died within 5 years after a surgical operation (5D group) and genes from patients survived free of disease for several years or more after the operation (5S group),
5 depending on their expression functions, in primary breast cancer.

[0021]

The gene correlated with prediction of the postoperative prognosis of breast cancer of the present invention is obtained by evaluating the data of a cDNA microarray using a Random-permutation test and a Mann-Whitney test. The present invention presents an
10 approach more useful at clinical level, by evaluating gene expression functions by a combination of a cDNA microarray and a semi-quantitative PCR experiment.

[0022]

In the present invention, a gene correlated with prediction of the postoperative prognosis of primary breast cancer has been identified by evaluating gene expression
15 functions in breast cancer patients.

Specifically, one aspect of the present invention is a gene selected from the following sequences selected from known sequences correlated with prediction of the postoperative prognosis of primary breast cancer;
pro-alpha-1 type 3 collagen (PIIIP),
20 complement component C1r,
dihydropyrimidinase-like 3 (DPYSL3),
protein tyrosine kinase 9-like (PTK9L),
carboxypeptidase E (CPE),
alpha-tubulin,
25 beta-tubulin,
heat shock protein HSP 90-alpha gene,
malate dehydrogenase,
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (NDUFB3).

[0023]

Some of the above-mentioned genes are believed to be correlated with proliferation or distant metastasis of tumor cells, and for example, a heat shock protein HSP 90-alpha is a chaperone for a lot of kinases, and has a possibility of promoting growth of cancer cells (Neckers, L. (2002) Trends Mol Med 8, S55-61). Malate dehydrogenase is an important enzyme correlated with energy accompanying aerobic or anaerobic metabolism, and the activity of malate dehydrogenase is correlated with a tumor marker for squamous cell carcinoma (Ross, C.D., et al. (2000) Otolaryngol Head Neck Surg 122, 195-200). NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3(NDUFB3) belongs to an mitochondrial electron transport chain, and chromosome abnormality in a region containing NDUFB3 is remarkable in a breast cancer cell line MDA-MB-231 (Xie, D., et al. (2002) Int J Oncol 21, 499-507).

[0024]

The above-mentioned 10 genes correlated with prediction of the postoperative prognosis of primary breast cancer show different expressions in a group of good prognosis (5S group) and a group of bad prognosis (5Y group), and 7 genes among the 10 genes are genes highly expressed in a group of good prognosis (5S group).

Namely, one aspect of the present invention is a gene selected from the following sequences highly expressed in a group of good prognosis selected from known sequences correlated with prediction of the postoperative prognosis of primary breast cancer;

pro-alpha-1 type 3 collagen (PIIIP),
complement component C1r,
dihydropyrimidinase-like 3 (DPYSL3),
protein tyrosine kinase 9-like (PTK9L),
carboxypeptidase E (CPE),
alpha-tubulin,
beta-tubulin.

[0025]

3 genes among the 10 genes correlated with prediction of the postoperative prognosis of primary breast cancer are genes highly expressed in a group of bad prognosis (5Y group). Namely, one aspect of the present invention is a gene selected from the following sequences highly expressed in a group of bad prognosis selected from known sequences correlated with prediction of the postoperative prognosis of primary breast cancer;
heat shock protein HSP 90-alpha gene,
malate dehydrogenase,
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (NDUFB3).

[0026]

Here, the prediction index (PI) for primary breast cancer is defined as described below and can be used for prediction of the postoperative prognosis of breast cancer.

Prediction index (PI) = (total of normalized expression ratios of the above-mentioned 7 genes highly expressed in a group of good prognosis in breast cancer tissue) - (total of normalized expression ratios of the above-mentioned 3 genes highly expressed in a group of bad prognosis in breast cancer tissue)

[0027]

In the present invention, gene expression functions in breast cancer patients have been evaluated and 10 genes correlated with prediction of the postoperative prognosis of node-negative breast cancer have been identified.

Specifically, one aspect of the present invention is a gene selected from the following sequences selected from known sequences correlated with prediction of the postoperative prognosis, in (node-negative)(n0) breast cancer with no metastasis to a lymph node in operation;

AF058701/ DNA polymerase zeta catalytic subunit (REV3),

AI066764/ lectin, galactoside-binding, soluble, 1 (galectin 1),

x15940/ ribosomal protein L31.,

Hs.94653/ neurochondrin (KIAA0607),

M13436/ ovarian beta-A-inhibin,

Hs.5002/ copper chaperone for superoxide dismutase; CCS,
D67025/ proteasome (prosome, macropain) 26S subunit, non-ATPase, 3,
M80469/ MHC class I HLA-J gene,
Hs.4864/ ESTs,
5 Hs.106326/ ESTs.

[0028]

The above-mentioned genes correlated with prediction of the postoperative prognosis of node-negative breast cancer include genes correlated with proliferation and distant metastasis of tumor cells. For example, galectin 1 is an autocrine type cancer repressor for regulating cell differentiation (AxelH, et al. (2003) Int. J. Cancer, 103: 370-379). Further,
10 a gene activating cancer metastasis is included.

[0029]

The above-mentioned 10 genes correlated with prediction of the postoperative prognosis of node-negative breast cancer show different expressions in a group of good
15 prognosis (5Y-F group) and a group of bad prognosis (5Y-R group), and 3 genes among the 10 genes are genes highly expressed in a group of bad prognosis (5Y-R group). Namely, one aspect of the present invention is a gene selected from the following sequences highly expressed in a group of bad prognosis selected from known sequences correlated with prediction of the postoperative prognosis, in node-negative breast cancer in operation;

20 AF058701/ DNA polymerase zeta catalytic subunit (REV3),
AI066764/ lectin, galactoside-binding, soluble, 1 (galectin 1),
x15940/ ribosomal protein L31.

[0030]

7 genes among the 10 genes correlated with prediction of the postoperative prognosis
25 of node-negative breast cancer are genes highly expressed in a group of good prognosis (5Y-F group). Namely, one aspect of the present invention is a gene selected from the following sequences highly expressed in a group of good prognosis selected from known sequences correlated with prediction of the postoperative prognosis, in node-negative breast

cancer;

Hs.94653/ neurochondrin (KIAA0607),

M13436/ ovarian beta-A-inhibin,

Hs.5002/ copper chaperone for superoxide dismutase; CCS,

5 D67025/ proteasome (prosome, macropain) 26S subunit, non-ATPase, 3,

M80469/ MHC class I HLA-J gene,

Hs.4864/ ESTs,

Hs.106326/ ESTs.

[0031]

10 Here, the prognosis score (PS) for node-negative breast cancer is defined as described below and can be used for prediction of the postoperative prognosis of breast cancer.

Prognosis score (PS) = (total of normalized expression ratios of the above-mentioned 3 genes highly expressed in a group of bad prognosis in breast cancer tissue) - (total of normalized expression ratios of the above-mentioned 7 genes highly expressed in a group of
15 good prognosis in breast cancer tissue).

[0032]

In the present invention, 20 genes correlated with prediction of the postoperative prognosis of estrogen receptor-negative breast cancer have been identified, by evaluating gene expression functions in breast cancer patients.

20 Specifically, one aspect of the present invention is a gene selected from the following sequences selected from known sequences correlated with prediction of the postoperative prognosis, in estrogen receptor-negative breast cancer;

Hs.108504/ FLJ20113/ ubiquitin-specific protease otubain 1

Hs.146550/ MYH9/ myosin, heavy polypeptide 9, non-muscle

25 Hs.194691/ RAI3/ retinoic acid induced 3

Hs.1975/ TDRD3/ tudor domain containing 3

Hs.203952/ TRRAP/ transformation/transcription domain-associated protein

Hs.278607/ GSA7/ ubiquitin activating enzyme E1-like protein

- Hs.429/ ATP5G3/ ATP synthase, H⁺ transporting,
mitochondrial F₀ complex, subunit c (subunit 9) isoform 3
Hs.75305/ AIP/ aryl hydrocarbon receptor interacting protein
Hs.81170/ PIM1/ pim-1 oncogene
- 5 Hs.99987/ ERCC2/
excision repair cross-complementing rodent repair deficiency, complementation group 2
Y12781/ Transducin (beta) like 1 protein
Hs.104417/ KIAA1205 protein
cl.21783/ Hypothetical protein
- 10 Hs.112628/ Hypothetical protein: MGC43581
Hs.170345/ Hypothetical protein FLJ13710
Hs.53996/ weakly similar to zinc finger protein 135
Hs.55422/ Hypothetical protein
Hs.112718/ EST
- 15 Hs.115880/ EST
Hs.126495/ EST
[0033]

The above-mentioned genes correlated with prediction of the postoperative prognosis of estrogen receptor-negative breast cancer include genes correlated with proliferation and distant metastasis of tumor cells. For example, PIM1 is serine/threonine kinase, and there is a correlation between clinical results of prostate cancer and the expression (Oesterreich, S., et al. (1996) Clin Cancer Res, 2, 1199-1206). TRRAP protein is a subunit of a mammal HTA complex, and antisense RNA against TRRAP inhibits estrogen-dependent growth of breast cancer cells.

25 [0034]

The above-mentioned 20 genes correlated with prediction of the postoperative prognosis of estrogen receptor-negative breast cancer show high expression in a group of bad prognosis (5y-D group). Namely, one aspect of the present invention is a gene

selected from known sequences correlated with prediction of the postoperative prognosis, in the above-mentioned estrogen receptor-negative breast cancer highly expressed in a group of bad prognosis.

[0035]

5 Here, postoperative prognosis of breast cancer can be predicted as described below, based on the expression of the above-mentioned gene correlated with prediction of the postoperative prognosis of estrogen receptor-negative breast cancer;

(1) when the expression levels in breast cancer tissue of the above-mentioned 20 genes correlated with prediction of the postoperative prognosis of estrogen receptor-negative
10 breast cancer are compared with the average value in a parent population, and if the expression level of each gene is 2-fold or more of the average value in a parent population, one point is imparted,

(2) when the procedure of (1) is carried out on 20 genes, and if the total point is 8 points or more, prognosis is decided to be bad.

15 [0036]

The above-mentioned gene correlated with prediction of the postoperative prognosis of breast cancer can be used as a marker for inspection of breast cancer postoperative prognosis. Namely, one aspect of the present invention is a method of inspecting the postoperative prognosis of breast cancer using the above-mentioned gene as a marker.

20 [0037]

The above-mentioned gene correlated with prediction of the postoperative prognosis of breast cancer can be used as a marker for screening of cancer therapeutic medicines for controlling the postoperative prognosis of breast cancer. Namely, one aspect of the present invention is a method of screening cancer therapeutic medicines for controlling the
25 postoperative prognosis of breast cancer using the above-mentioned gene as a marker.

[0038]

The above-mentioned gene correlated with prediction of the postoperative prognosis of breast cancer can be used as a marker for diagnosis of the postoperative prognosis of breast

cancer. It is also possible to design probes specific to the above-mentioned gene and to use these probes as a marker. These probes can be designed, for example, by Probe Quest (registered trademark) manufactured by Dyna Com. Namely, one aspect of the present invention is a diagnosis kit for the postoperative prognosis of breast cancer containing a reagent using the above-mentioned gene as a marker.

[0039]

The above-mentioned diagnosis kit can include a microarray. Namely, one aspect of the present invention is the diagnosis kit, wherein the diagnosis kit includes a microarray.

[0040]

The microarray of the above-mentioned diagnosis kit including a microarray includes a fiber type microarray. Here, for a method of preparing a fiber type microarray, the above-mentioned Patent Literatures 6-7 are cited. Namely, one aspect of the present invention is the above-mentioned diagnosis kit wherein the microarray is a fiber type microarray.

[0041]

Next, aspects of the present invention will be specifically illustrated by examples, but the present invention is not limited to these examples.

[Example 1]

[0042]

Evaluation of gene expression function for prediction of the postoperative prognosis in estrogen receptor-negative breast cancer

[0043]

(Tissue sample)

An informed consent was obtained according to a guide line accepted by an ethics committee of Cancer Society and by Nippon Medical School, then, primary breast cancer and tissue from adjacent normal mammary gland were collected from breast cancer patients who undergone an operation in 1995 to 1997 in Cancer Society attached hospital (Tokyo). The tissue was quickly frozen and preserved at -80°C. For 954 patients, all members were

clinically traced during a period of 5 years or more or until death, and samples were selected from 10 estrogen receptor-negative breast cancer patients died within 5 years after the operation (5y-D) and 10 patients survived free of disease for 5 years or more after the operation (5y-S). The backgrounds of both the patient groups were allowed to coincide in age, lymph node metastasis, tumor diameter and tissue type (Table 1).

[0044]

(Clinical feature of 20 cases of breast cancer)

[0045]

[Table 1]

group	Case No.	ER condition	Age	Sex	Process ^a	TNM classification ^b		TTD ^c
						Tumor	Lymph node	
5y-D	3281	Negative	34	Female	a2	T2	N1b	9
	3459	Negative	64	Female	a2	T4	N3	6
	3550	Negative	73	Female	a2	T4	N1b	12
	3892	Negative	62	Female	a2	T2	N1a	21
	3948	Negative	60	Female	a2	T2	N1a	51
	4020	Negative	50	Female	a2	T2	N3	28
	3654	Negative	46	Female	a2	T4	N1b	19
	4118	Negative	53	Female	a2	T1	N1a	21
	4462	Negative	34	Female	a1	T2	N1a	24
	4126	Negative	51	Female	b5	T4	N3	6
5y-S	3656	Negative	31	Female	a2	T2	N1a	>60
	3197	Negative	42	Female	a1	T1	N1a	>60
	3662	Negative	58	Female	a2	T2	N0	>60
	3241	Negative	47	Female	a2	T2	N1a	>60
	3267	Negative	51	Female	a2	T2	N1a	>60
	3329	Negative	60	Female	a2	T2	N1a	>60
	3345	Negative	43	Female	a1	T2	N2	>60
	3556	Negative	59	Female	a2	T3	N0	>60
	3558	Negative	57	Female	a2	T3	N1b	>60
	3658	Negative	42	Female	a1	T2	N1a	>60

^aa1: invasive papillotubular carcinoma. a2: invasive solid-tubular carcinoma. b5: squamous cell carcinoma.

^bTNM classification: clinical classification by Japan Breast Cancer Society

^cTTD: time to death after surgery (months)

10

[0046]

All patients underwent postoperative adjuvant therapy according to “Postoperative clinical protocol for breast cancer (nyugan no tameno shujutsugo no rinsho no purotokoru)” of Cancer Society attached hospital. In each case, selection of adjuvant therapy was determined strictly based on surgical operation type, lymph node involvement condition, and presence of local or distant metastasis. In the study of the present invention, all patients did not have distant metastasis before the adjuvant chemical therapy and did not undergo radiation therapy or chemical therapy before the surgical operation.

[0047]

(Clinicopathological parameter)

The following parameters were checked: tissue type, tumor diameter and invasion (t factor), lymph node involvement, and conditions of estrogen receptor (ER) and progesterone receptor (PgR). Tumors were classified into the following types according to TNM classification and to tissue classification of Japan Breast Cancer Society (1989); noninvasivetubular (1a), invasivepapillotubular (a1), invasive solid-tubular (a2), invasivescirrhoucarcinoma (a3), and other special types (b). The classification is basically the same as breast cancer tissue classification of WHO. t factors were classified into the following types according to histological TNM classification; tumor with a maximum size of 2 cm or less (t1), tumor with no invasion into skin or pectoral muscle and with a maximum size of 2 cm or more (t2), and tumor with invasion into skin or pectoral muscle (t3).

[0048]

15 (Design and construction of cDNA microarray)

From 25344 cDNAs selected from UniGene database, “genome wide cDNA microarray” was constructed. The cDNAs were made by RT-PCR using poly(A)+RNAs separated from various human organs. The PCR products were spotted on slide glasses of type 7 (Amersham Biosciences UK Limited, Buckinghamshire, UK) using Array Spotter Generation III (Amersham Biosciences). Each slide contains 384 house-keeping genes.

[0049]

(Preparation and proliferation of RNA)

A tumor raw material was quickly frozen at -80°C immediately after collection. RNA was extracted using TRIzol (Invitrogen Inc., Carlsbad, CA, USA), further, purified using RNeasykits (Quiagen Inc., Valencia, CA). The purity of each RNA was evaluated by a spectrophotometry and electrophoresis on 1.2% modified formamide gel. The high purity RNA was defined as a sample having an absorbance ratio (260 nm/280 nm) of 1.8 to 2.0 and in which 28S/18S liposomal bands show a ratio of 1.8 or more on formamide gel

electrophoresis. After treating with 1 unit of DNaseI (Epicentre Technologies, Madison, WI)(1 unit/ μ l), RNA amplification by T7RNA polymerase was carried out using 2 μ g of RNA from each sample as a starting raw material. Amplification was carried out twice, and the amplified RNA (aRNA) was purified by RNeasykits (Quiagen Inc., Valencia, CA).

- 5 The amount of each aRNA was measured by a spectrophotometer, and the quality was checked by formamide gel electrophoresis.

[0050]

(Labeling of aRNA, hybridization and scanning)

- cDNA for microarray analysis was prepared from aRNA. aRNAs (5 to 10 μ g) from
10 breast cancer and normal mammary gland tissue were labeled with Cy5 (cancer sample) and Cy3 (normal sample) using aminoallyl-cDNA labelingkits (Ambion, Austin, TX). The Cy3-labeled cDNA probe and the Cy5-labeled cDNA probe were mixed and heated at 95°C for 5 minutes, then, quenched with ice for 30 seconds, and hybridized on a microarray. The mixed probes were added to formamide (Sigma-Aldrich Corp., St. Louis, MO, USA)
15 having a 50% final concentration of microarrayhybridization solution version 2 (Amersham Biosciences UK Limited, Buckinghamshire, UK). After hybridization at 40°C for 15 hours, the microarray slides were washed first with 1xSSC and 0.2% SDS at 55°C for 10 minutes, then, washed twice with 0.1xSSC/0.2% SDS each for 1 minute at room temperature. All treatments were carried out by Automated Slide Processor System (Amersham). The
20 signal strength of each hybridization was scanned by Gene Pix 4000A (Axon Instruments, Inc., Foster City, CA, USA), and evaluated by Gene Pix 3.0 (Axon Instruments) by a spectrophotometry. The scanned signals were normalized by a method described in the following literature (the total gene normalization method) (Yang YH, Dudoit S, Luu P, et al. (2002) Nucleic Acids Res 30, e15; Manos EJ, Jones DA. (2001) Cancer Res 61: 433-348).

25 [0051]

(Signal analysis and selection of genes showing different expressions)

The signal strength of each hybridization was evaluated by a photometry by Gene Pix 3.0 (Axon Instruments, Inc., Foster City, CA, USA). For normalizing mRNA expression

levels between cancer and control, the Cy5: Cy3 ratio in each gene expression was adjusted. As a result, the averaged Log (Cy5: Cy3 ratio) of the house keeping genes was zero. 27 house keeping genes were adopted from a house-keeping panel in Web site <http://www.nhgri.nih.gov/DIR/LCG/ARRAY/expn.html>. For each microarray slide, the cut off value of (S/N) ratio was set at 3.0. Genes with signal strengths of Cy3 and Cy5 lower than the cut off value were excluded out of the investigation.

[0052]

(Mann-Whitney test)

For investigating genes showing apparently different expressions between 5y-D tumor and 5y-S tumor, Mann-Whitney test was applied to a series of samples X. X represents Cy5/Cy3 signal strength ratio of each gene and each sample (OnoK, Tanaka T, Tsunoda T, et al. (2000) Cancer Res 2000; 60: 5007-5011). The U value was calculated for genes imparting significant signals in at least 5 samples in both groups. Genes showing U values of lower than 23 or larger than 77 were selected. Since the U value is obtained by calculation for 5y-S group based on 5y-D group in each gene based on each X value, U values lower than 23 were evaluated to manifest higher expression in 5y-S group than in 5y-D group. However, genes with U values higher than 77 were evaluated to manifest higher expression in 5y-D group than in 5y-S group. Base on this criterion, 183 genes were highly expressed in 5y-S group and 31 genes were highly expressed in 5y-S group. Thus, only genes in which intermediate expression values show a difference of 2-fold or more between two groups ($\mu_{XD}/\mu_{XS} \leq 0.5$ or ≥ 2.0 , μ_{XD} and μ_{XS} represent average X values in 5y-D and 5y-S group, respectively) were defined as genes correlated with prognosis. As a result, 110 genes in total were selected. Of them, 90 genes were expressed at higher level in 5y-D tumor group and 20 genes were expressed at higher level in 5y-S tumor group.

[0053]

(Random-permutation test)

Further, for evaluating values of 110 genes selected by the Mann-Whitney test, a

permutation test was carried out. A possibility, P_s of a gene for correlating with a group difference was also assumed. When each gene is represented by an expression vector $v(g) = (X_1, X_2, \dots, X_{20})$ (X_i shows a gene expression level of i -th sample in the first sample group), an ideal expression pattern is expressed by $c = (c_1, c_2, \dots, c_{20})$ ($c_i = +1$ or 0 , depending on whether i -th sample belongs to S group or D group).

Correlation between a gene and a group difference P_{gc} was defined as described below. That is, $P_{gc} = (\mu_S - \mu_D) / (\delta_S + \delta_D)$; $\mu_S(\mu_D)$ and $\delta_S(\delta_D)$ show standard deviation of $\log_2 X$ of the gene “g” of each sample in a newly defined S (or D) group.

The permutation test was carried out while substituting the coordinate of c . The correlation values, P_{gc} were calculated between all permutations. These procedures were repeated for 10000 times. Accidentally, the p value showing a possibility of a gene for classifying two groups was evaluated for each of 110 genes selected. Finally, 71 gene highly expressed in 5y-D case and 15 gene expressed low in 5y-S case were selected.

[0054]

15 (Semi-quantitative RT-PCR)

RNA (2 μ g) was treated with DNase I (Epicentre Technologies, Madison, WI, USA), and single-stranded cDNAs were subjected to reverse transcription using Reverscript II reverse transcriptase (manufactured by Wako Pure Chemical Industries, Ltd., Osaka, Japan) and oligo (dT) 12-18 primer. Single-stranded cDNAs were adjusted in the concentration for the subsequent PCR amplification by monitoring expression of GAPD (glyceraldehyde-3-phosphatedehydrogenase) as a quantitative control. Each PCR was carried out under the following reaction conditions using Gene Amp PCR system 9700 (Applied Biosystems, Foster City, CA, USA) at an amount of 1xPCR buffer of 30 μ l.

94°C 5 minutes,

25 (94°C 30 seconds, 60°C 30 seconds, and 72°C 30 seconds) for 25 to 35 cycles.

Primer sequences used in RT-PCR are as described below:

SEQ ID No. 1 GAPD (control) forward, 5'-GGA AGGTGA AGG TCG GAG T-3'

SEQ ID No. 2 reverse, 5'-TGG GTG GAA TCA TAT TGGAA-3';

SEQ ID No. 3 Hs.108504F, 5'-ACA CTT CAT CTG CTCCT CAT AG-3';

SEQ ID No. 4 Hs.108504R, 5'CTG CCT AGA CCT GAGGAC TGT AG-3';

SEQ ID No. 5 Hs.146550F, 5'ACT GAG GCC TTT TGGTAG TCG-3';

SEQ ID No. 6 Hs.146550R, 5'TCT CTT TAT TGT GATGCT CAG TGG-3';

5 SEQ ID No. 7 Hs.76607F, 5'AAA TCC TTC TCG TGT GTTGAC TG-3';

SEQ ID No. 8 Hs.76607R, 5'CAG TCA TGA GGG CTA AAAACT GA-3';

SEQ ID No. 9 Hs.1975F, 5'GAA GAC AAC AAG TTT TAC CGG G-3';

SEQ ID No. 10 Hs.1975R, 5'ATG GTT TTA TTG ACG GCAGAA G-3';

SEQ ID No. 11 Hs.203952F, 5'AGG ACA CGT CCT CTCCTC TCT C-3';

10 SEQ ID No. 12 Hs.203952R, 5'TAA AGC TAG CGA AGGAAC GTA CA-3';

SEQ ID No. 13 Hs.278607F, 5'TCC CTT CTG TTT CCT CAG TGT T-3';

SEQ ID No. 14 Hs.278607R, 5'CCT GCC CCG ATA AAA ATA TCT AC -3';

SEQ ID No. 15 Hs.429F, 5'TTG ACC TTA AGC CTC TTTTCC TC-3';

SEQ ID No. 16 Hs.429R, 5'ATA ACG TAC ATT CCC ATGACA CC-3';

15 SEQ ID No. 17 Hs.75305F, 5'ACT TTC AAG ATG GGACCA AGG-3';

SEQ ID No. 18 Hs.75305R, 5'ATA TAC ACA GAA GCATGA CGC AG-3';

SEQ ID No. 19 Hs.81170F, 5'TTG CTG GAC TCT GAAATA TCC C-3';

SEQ ID No. 20 Hs.81170R, 5'TTC CCC TGT ACA GTATTT CAC TCA-3';

SEQ ID No. 21 Hs.99987F, 5'CTG AGC AAT CTG CTCTAT CCT CT-3';

20 SEQ ID No. 22 Hs.99987R, 5'GTT CCA GAT TCG TGAGAA TGA CT-3';

SEQ ID No. 23 Y12781F, 5'ACC AGT AAC AAC TGT GGGATG G-3';

SEQ ID No. 24 Y12781R, 5'CAA ATG AGC TAC AAC ACACAA GG-3';

SEQ ID No. 25 Hs.104417F, 5'CCC CCT CCA CCT TGTACA TAA T-3';

SEQ ID No. 26 Hs.104417R, 5'GTT TTC GTT TGG CTGGTT GTG-3';

25 SEQ ID No. 27 cl.21783F, 5'GTC TGA GAT TTT ACTGCA CCG-3';

SEQ ID No. 28 cl.21783R, 5'GGA TGG AGC TGG AGGATA TTA-3';

SEQ ID No. 29 Hs.112628F, 5'ATT GCT AAG GAT AAGTGC TGC TC-3';

SEQ ID No. 30 Hs.112628R, 5'TGT CAG TAT AGA AGCCTG TGG GT-3';

SEQ ID No. 31 Hs.170345F, 5'TTC TTA GGC CAT CCCTTT TCT AC-3';
 SEQ ID No. 32 Hs.170345R, 5'GCA TCT GAA TGT CTTTCT CCC TA-3';
 SEQ ID No. 33 Hs.53996F, 5'CCA TAG GAT CTT GACTCC AAC AG-3';
 SEQ ID No. 34 Hs.53996R, 5'ACT GGG AGT GGA GGAAAT TAG AG-3';
 5 SEQ ID No. 35 Hs.55422F, 5'CTA ATG TAA GCT CCATTG GGA TG-3';
 SEQ ID No. 36 Hs.55422R, 5'CAA ACT GCA AAC TAGCTC CCT AA-3';
 SEQ ID No. 37 Hs.112718F, 5'AAG ACT AAG AGG GAA AAT GTG GG-3';
 SEQ ID No. 38 Hs.112718R, 5'AGG TAA CCC AAA GTG ACA AAC CT-3';
 SEQ ID No. 39 Hs.115880F, 5'TTA AGT GAG TCT CCT TGG CTG AG-3';
 10 SEQ ID No. 40 Hs.115880R, 5'AGG GCC CCT ATA TCC AAT ACC TA-3';
 SEQ ID No. 41 Hs.126495F, 5'GAT CTT TCA AGA TGAGCC AAG GT-3';
 SEQ ID No. 42 Hs.126495R, 5'AGT CAT TCA GAA GCCATT GAG AC-3'
 [0055]

(Measurement of signal strength of RT-PCR product and calculation of prognosis score)

15 A PCR product was detected by 2% agarose gel electrophoresis and ethidium bromide staining. A gel was scanned by a digital image processing system (AlphaImager 3300; Alpha Innotech, San Leandro, CA, USA) according to the Spot Density method. A two-dimensional region of each band was constructed, and pixel strength (gene expression) was obtained in which the density was defined as IDV (Integrated Density Value).
 20 Importance in a difference in IDV in each group was evaluated by the Student's t-test. As a result, 20 genes showing p values of 0.05 or lower in the t-test were selected as a candidate (Table 2). That is, expression levels of the 20 genes were significantly higher in the 5y-D group than in the 5y-S group. Base on this information, the present inventors have tried to establish a scoring system for predicting the postoperative prognosis. In this procedure,
 25 each gene was determined depending on whether the expression level of each sample was higher than the average expression level of 20 samples or not. When the expression level of a sample was 2-fold or more than the average, + 1 point was imparted additionally. Next, points of all of the 20 genes were summed up for obtaining the total vote (prognosis

score) for each sample. As a result, a case of a sample of 8 points or more was evaluated as an indication of bad prognosis. On the other hand, a case of a sample of 8 points or less was evaluated as an indication of preferable prognosis.

[0056]

(20 candidate genes of prognosis scoring system)

[0057]

[Table 2]

Hs./Accession No.	kind
Hs.108804	FLJ20113: ubiquitin-specific protease otubain 1
Hs.148550	MYH9: myosin, heavy polypeptide 9, non-muscle
Hs.194691	RAI3: retinoic acid induced 3
Hs.1975	TD RD3: tudor domain containing 3
Hs.203952	TRRAP: transformation/transcription domain-associated protein
Hs.278607	GSA7: ubiquitin activating enzyme E1-like protein
Hs.429	ATP5G3: ATP synthase, H+ transporting, mitochondrial F0 complex, subunit c (subunit 9) isoform 3
Hs.75305	AIP: aryl hydrocarbon receptor interacting protein
Hs.81170	PIM1: pim-1 oncogene
Hs.99987	ERCC2: excision repair cross-complementing rodent repair deficiency, complementation group 2
Y12781	Transducin (beta) like 1 protein
Hs.104417	KIAA1205 protein
cl.21783	Hypothetical protein
Hs.112628	Hypothetical protein: MGC43581
Hs.170345	Hypothetical protein FLJ13710
Hs.53996	weakly similar to zinc finger protein 135
Hs.55422	Hypothetical protein
Hs.112718	EST
Hs.115880	EST
Hs.126495	EST

5 [0058]

(Result)

257 genes highly expressed significantly in estrogen receptor-negative breast cancer tissue were clarified, and 378 genes expressed low were clarified likewise. For identifying genes showing different expressions between the 5y-D group and the 5y-S group, the data of a microarray was analyzed by the Mann-Whitney test and the Random-permutation test. As a result, 71 genes in total (including 10 EST and 9 genes encoding virtual protein) in 5y-D tumor were classified in common into a group of higher expression. In contrast, 15 genes (including 3 EST) were classified in common into a group of lower expression (Fig. 1).

15 [0059]

Genes highly expressed in the 5y-D group include the following genes correlated with proliferation and metastasis of cancer cells; matrix metalloproteinase 2 (MMP2), heat shock protein 27 HSPB1), Pim-1 oncogene (PIM1) and transformation/transcription domain-associated protein (TRRAP).

20 Genes expressed low in the 5y-D group include genes of HLA-C (major histocompatibility complex, class I, C) and specific kinase. A lot of genes having

correlations with DNA repair, transcription, signal transduction, cytoskeleton and adhesiveness showed different expressions between two groups.

[0060]

For confirming reliability of the data of a microarray, 20 genes highly expressed in the 5y-D group were selected (Hs.108504, Hs.146550, Hs.194691, Hs.1975, Hs.203952, Hs.278607, Hs.429, Hs.75305, Hs.81170, Hs.99987, Y12781, Hs.104417, cl.21783, Hs.112628, Hs.170345, Hs.53996, Hs.55422, Hs.112718, Hs.115880, and Hs.126495), and the expression levels of the genes were checked by semi-quantitative RT-PCR. The result coincided with the data of a microarray, and had a statistical significance for classifying the 5y-D group and the 5y-S group (typical data is shown in Fig. 2).

[0061]

For constructing a scoring system for predicting the postoperative prognosis using the expression profile of a marker gene, prognosis score was calculated by the above-mentioned method. Briefly, a marker gene was selected according to the following standard.

- (1) Higher signal strength than cut off level is shown in at least 60% of cases checked;
- (2) $|\mu_D - \mu_S|$ is 1.0 or less. Here, $\mu_D(\mu_S)$ shows an average value derived from logarithm converted relative expression ratio in the case of 5y-D(5y-S).

[0062]

Next, for identifying a marker gene capable of classifying the 5y-D group and the 5y-S group depending on the expression function, the Mann-Whitney test and the Random-permutation test were carried out. The result of a microarray correlated was confirmed by a semi-quantitative RT-PCR experiment. By the Student's t-test, 20 genes were selected as a prognosis marker (Table 2).

[0063]

Depending on the prognosis score (PS) of the present invention, 20 patients were divided into 10 members predicted to show poor prognosis (PS is 11 or more) and 10 members predicted to show excellent prognosis (PS is less than 11). As a result, it was shown by comparison with the postoperative progress which the scoring system of the

present invention has reliability with an accuracy of 80% in the 5y-D case and with an accuracy of 100% in the 5y-S case (Fig. 3A).

[0064]

5 Using the prognosis scoring system of the present invention, additional 5 cases were checked (Fig. 3B). The system predicted poor prognosis in 2 cases (PS > 11; patient TD-1 and patient TD-2), and excellent prognosis in 3 cases (PS < 11; patients TD-3, TS-1 and TS-2). As a result, this scoring system showed an accuracy of 80% regarding actual clinical results of these 5 cases.

[Example 2]

10 [0065]

Evaluation of gene expression function for prediction of the postoperative prognosis in node-negative breast cancer

[0066]

(Tissue sample)

15 A tissue sample was collected in the same manner as described in Example 1. Gene expression was investigated for tumors from 12 patients of node-negative (n0) cancer showed recurrence within 5 years after an operation (5Y-R) and 12 patients survived free of disease for 5 years or more after the operation (5Y-F). The clinical backgrounds of both the patient groups were allowed to coincide in age, lymph node metastasis, tumor diameter,
20 condition of hormone receptor, and pathological tissue (Table 3). The follow up intermediate period was 7.8 years, and the average period between the initial operation and recurring was 2.7 years in the 5Y-R group. All patients underwent the adjuvant therapy described in Example 1.

[0067]

25

(Clinical pathological data)

[0068]

[Table 3]

Case	Age	Climacteric condition	Histological classification ^a	Position	Diameter (mm)	TNM classification ^b			Stage	ER(+/-)	PgR(+/-)	D.F.I. ^c
						T	N	M				
R-1	55	Post.	a2	Rt	25	2	1a	0	II	+	-	12m
R-2	50	Pre.	a3	Lt	25	2	1a	0	II	+	+	16m
R-3	42	Pre.	a2	Rt	25	2	0	0	II	+	+	49m
R-4	39	Pre.	a3	Rt	35	2	0	0	II	+	-	20m
R-5	38	Pre.	a2	Lt	30	2	0	0	II	+	+	52m
R-6	61	Post.	a3	Lt	34	2	0	0	II	-	-	14m
R-7	54	Post.	b3	Lt	30	2	0	0	II	-	-	24m
R-8	37	Pre.	a2	Rt	23	2	0	0	II	-	-	25m
R-9	54	Post.	a3	Lt	25	2	1a	0	II	+	+	47m
R-10	83	Post.	a2	Rt	28	2	1a	0	II	+	+	38m
R-11	62	Post.	a2	Lt	23	2	0	0	II	-	+	40m
R-12	50	Post.	a3	Lt	35	2	0	0	II	-	-	25m
F-1	48	Pre.	a2	Lt	18	2	0	0	II	+	+	8Y
F-2	62	Post.	a2	Rt	25	2	0	0	II	+	-	8Y
F-3	57	Post.	a1	Rt	20	1	0	0	I	+	+	7Y10m
F-4	61	Post.	a2	Lt	30	2	1a	0	II	-	-	7Y2m
F-5	42	Pre.	a1	Lt	12	1	1a	0	I	-	+	7Y11m
F-6	51	Pre.	a2	Rt	28	2	1a	0	II	-	-	7Y10m
F-7	59	Post.	a2	Rt	40	3	0	0	II	-	-	7Y5m
F-8	57	Post.*	a2	Rt	45	3	1b	0	II	-	-	7Y5m
F-9	42	Pre.	a1	Lt	48	2	1a	0	II	-	+	7Y3m
F-10	58	Post.	a2	Lt	13	2	0	0	II	-	-	7Y3m
F-11	50	Post.	a2	Lt	25	2	0	0	II	+	+	7Y8m
F-12	55	Post.	a1	Rt	35	2	0	0	II	+	+	7Y5m

^a a1 : invasive papillotubular carcinoma, a2 : invasivesolid-tubularcarcinoma, a3 : invasive schirrhous carcinoma

^b TNM classification: clinically classified according to TNM classification by Japan Breast Cancer Society

^c D.F.I.: period of no pathogeny (disease free interval)

[0069]

(Clinicopathological parameter)

The clinicopathological parameter was checked by the method described in Example 1. The histological grade was evaluated by a method of Elastonand Ellis (Abrams JS. Breast Cancer 2001; 8: 298-304). Lymphoduct invasion was evaluated to be deficient or positive (for example, evaluated to be positive when one or more cancer cells are present in lymphoducts around cancer). Fatinvasion was evaluated to be deficient or positive (for example, evaluated to be positive in the case of invasion into interstitial tissue).

[0070]

(Preparation of cDNA microarray)

“Genome wide cDNA microarray kit (Amersham Biosciences UK Limited, Buckinghamshire, UK)” with 25344 cDNAs was used. The PCR product was stopped on
5 type 7 glass slides (Amersham Biosciences) using Array Spotter Generation III (Amersham Biosciences).

[0071]

(Preparation and proliferation of RNA)

Preparation and proliferation of RNA were carried out in the same method as
10 described in Example 1.

[0072]

(Labeling of aRNA, hybridization and scanning)

Labeling of aRNA, hybridization and scanning were carried out in the same method as described in Example 1.

15 [0073]

(Mann-Whitney test)

For identifying genes showing different expressions between a group of no disease and group of recurrence, normalized signals were analyzed by the Mann-Whitney test applied to a series of Xs. Here, X represents Cy5/Cy3 signal strength ratio of each gene and each
20 sample. Genes showing a difference of 2-fold or more in expression strength between two groups were selected. Genes with signal-noise ratios of 3.0 or less were excluded from analysis.

The U value was calculated for genes imparting significant signals in at least 5 samples in both groups. Genes with U values of lower than 37 or larger than 107 were
25 selected. Since the U value was obtained by calculation for 5Y-F group based on 5Y-R group in each gene based on each X value, genes with U values lower than 37 were evaluated to manifest higher expression in 5Y-F group than in 5Y-R group (first category). On the other hand, genes with U values higher than 107 were evaluated to manifest higher

expression in 5Y-R group than in 5Y-F group (second category).

Based on this method, 78 genes were identified in the first category and 55 genes were identified in the second category. Thus, only genes showing a difference of 2-fold or more of the intermediate expression value between two groups ($\mu X_R/\mu X_F \leq 0.5$ or ≥ 2.0 , μX_R and μX_F represent average X values in 5Y-R and 5Y-F group, respectively) were defined as genes correlated with prognosis. In total, 98 genes were selected, and of them, 64 genes showed higher expression level in 5Y-F tumor and 34 genes showed higher expression level in 5Y-R tumor.

[0074]

10 (Random-permutation test)

For evaluating values of genes selected by the Mann-Whitney test, a permutation test was carried out, and correlation to group difference (P_s) of genes selected was evaluated. When each gene is represented by an expression vector $v(g) = (X_1, X_2, \dots, X_{24})$ (X_i shows a gene expression level of i -th sample in the first sample set), an idealized expression pattern is expressed by $c = (c_1, c_2, \dots, c_{24})$ ($c_i = +1$ or 0 , depending on whether i -th sample belongs to F group or R group).

Correlation between a gene and a group difference P_{gc} was defined as described below. That is, $P_{gc} = (\mu_F + \mu_R)/(s_F + s_R)$; $\mu_F(\mu_R)$ and $s_F(s_R)$ show standard deviation of $\log_2 X$ of the gene “g” of each sample in a newly defined “F” group or “R” group.

20 The permutation test was carried out while substituting the coordinate of c . The correlation values, P_{gcs} were calculated between all permutations. These procedures were repeated for 10000 times. Accidentally, the p value showing a possibility of a gene for classifying two groups was evaluated for each of 58 genes selected.

[0075]

25 (Semi-quantitative RT-PCR)

RNA (5 μ g) was treated with DNase I (Epicentre Technologies, Madison, WI, USA), then, single-stranded cDNAs were subjected to reverse transcription using Reverscript II reversetranscriptase (manufactured by Wako Pure Chemical Industries, Ltd., Osaka, Japan)

and 0.5 µg/µl oligo (dT) 12-18 primer. The preparations of single-stranded cDNAs were diluted for the subsequent PCR amplification by monitoring GAPDH as a quantitative control. All PCRs were carried out under the following reaction conditions using Gene Amp PCR system 9700 (Applied Biosystems, Foster City, CA, USA) at an amount of

5 1xPCR buffer of 30 µl.

94°C 2 minutes,

(94°C 30 seconds, 58-62°C 30 seconds, and 72°C 30 seconds) for 27 to 35 cycles

72°C 5 minutes.

Primer sequences for RT-PCR of GAPDH are as described below:

10 SEQ ID No. 43 (forward) 5'-GAA AGG TGA AGG TCG GAG T-3'

SEQ ID No. 44 (reverse) 5'-TGG GTG GAA TCA TAT TGG AA-3'

[0076]

(Primer of semi-quantitative PCR (gene highly expressed in group of no disease))

[0077]

15 [Table 4A]

Ac./HS	SEQ ID No.	Forward	SEQ ID No.	Reverse
M90439	45	CCAGACATCCATGGTACCTATAA	46	TATGCATTGAAACCTTACAGGGG
AF047472	47	CTGTAAACAAAGCGAGGTTAAGG	48	GGGTTCTGCATCTCGTTTATTAG
Hs.118251	49	GACACATAGCTCATAGGCACACA	50	TTCTGGTACATGGTAAGTGCTCA
D26125	51	TCCGCCATATTGATTCTGCTTA	52	GTTTGCTTTCTGGACCATGGATA
Hs.8619	53	GATAACAACCTGGACCACATCCC	54	AACAGGCAGACGAGGTAGACAC
X16135	55	GAGAAGGATGGGTCCACCAGT	56	GTACATGGGCAGCACAAATGTAT
Hs.9006	57	ATTTCAATTGGTAGTATGGCCAC	58	ATACCATGGGACAGGATTGTAAG
M18963	59	GCTCAGACCAGCTCATACTTCAT	60	CCAAAGACTGGGGTAGGTAAGAC
X07979	61	CTGGTGCTTTCTATCACCTCTTC	62	GACTAGTGTGAAACAAGATGGGC
AF018080	63	CTTGAACCCAGGAGTTTGAGAC	64	GTGCCTCAGCTTTCTGAGTAGC
Hs.58464	65	CTGGTGCTGACTATCCAGTTGA	66	CTGGTAAACTGTCCAAAACAAGG
S79867	67	CTCTTACCTGGACAAGGTGCGT	68	GGATGAGCTCTGCTCCTTGAG
J02854	69	CAATGTTTGACCAAGTCCCAGA	70	CATGTTGTCTCAGTCTCTATTGG
Z35309	71	GGACAGCAGCTGGAGTACACA	72	AATCAGATTTGTCCGGTGCCTT
Hs.83097	73	GGCTCTGCACTAAGAACACAGAG	74	ACAACCTAGCTCTCAGTTCAGGCA
Hs.79137	75	TGGAGCAGTATGACAAGCTACAA	76	AAGCAGCACTGCATAAAGCTGTT
Hs.4864	77	TAACTACTTTCTGTGGTGCGCT	78	CCACAAACAGGAAGCTATGTTCT
Y00052	79	GTACTATTAGCCATGGTCAACCC	80	CTACAGAAGGAATGATCTGGTG
Hs.5002	81	ATCAGTACGGGGACCTTACAAAC	82	CCTGTACTGAGCTCTCCAAAGAC
U43519	83	TCCCTAGCTTCTCTCCACA	84	AGAATCATGCCTCCCCTTCT
Hs.94653	85	ACCCCTCAAGTGTAAGGAACTG	86	GGATCAAGAGTGTGTGTGTGTGT
X51441	87	CAATGCCAGAGAGAATATCCAGA	88	GATACCCATTGTGTACCCCTCTCC
Hs.108623	89	CCACTCCACATAAGGGGTTTAG	90	GAGGTTCTAGCTAAGTGCAGGGT
Hs.5318	91	CCATTGACATTGGAGTTAAGTATGC	92	GGCAAAGACCACATTTAGCAAT
Hs.69469	93	GAAAGCCTATGTGAAAAGCTGGT	94	TTGTTTCCAGGCATTAAGTGTG
AA777648	95	GCATCTTAGTCCACACAGTTGGT	96	GCCCTTACAGGTGGAGTATCTTC
Hs.106131	97	CTCATAGCCAGCATGACTTCTTT	98	GGTTCACTTGTGACTGGTCATCT
X54079	99	ACTTTTCTGAGCAGACGTCCAG	100	TATCAAAAGAACACACAGGTGGC
AJ041182	101	ACGTTATTCCCAGTTCCTAAACC	102	AGTCTCGGGTGACTCAATATGAA
AA148265	103	AGTTGAACCCAGGTACCTTTCTC	104	CTAGGCCCTTTTAGAAAACATGG
Hs.4943	105	TACTGGGAACGACTAAGGACTCA	106	TGCTGTGTTGAGTAGGTTTCTGA
Hs.106326	107	TGAGAGTCCTCAGAGGGTATCAG	108	CTTGAAGTCAAGAGTCCTGGTGT
M13436	109	TTTCTGTTGGCAAGTTGCTG	110	CCCTTTAAGCCCACTTCCTC
X99920	111	GATGAGAAGATGAAGAGCTTGGA	112	GAGGAAGCTTTATTTGGGAAGAG
U22970	113	ACTTCCCTCTCTGCCTTTCTG	114	CAGATTGTTTGGGCTTCTCACT

[0078]

(Primer of semi-quantitative PCR (gene highly expressed in group of recurrence))

5 [0079]

[Table 4B]

Ac./HS	SEQ ID No.	Forward	SEQ ID No.	Reverse
X75252	115	GTCTGGTCAGCTTTGCTTCC	116	GGCAAGTTCTGCACAGATGA
AA989127	117	CAGCTCAGTGCCACCATGAAT	118	GTGGGACTGAGATGCAGGAT
Hs.128520	119	CACGGACTCATGAATGTAGTGAA	120	GTGTAGTGGCAGCATCATAGCTT
HSMLN50	121	GGGACCAAACAGACCAAAGA	122	CACCCACAGAGCCTGTATT
AF058701	123	CGGAAAGGCACTATTTACAAT	124	ACAGGCCACAGGTTTGTAA
AF043473	125	AAGCTCTTCAGCTGCGTCTC	126	CCTCCTCCTTTTCAGCTGTG
Hs.26052	127	TCTGGAACCCTAAAAGTGTCTG	128	TCTTTCAACATCTCTCCACCCTA
Hs.77961	129	AGATACCTGGAGAACGGGAAG	130	GGAAAGTAAGAAGTTGCAGCTCAG
Hs.26484	131	ATTAGGTTTCACCCAAAG	132	AGACGAGACTTGTCTTCTC
U44798	133	CAGGGACTTGGTCACAGGTT	134	TTCTTCTCCCTCCCCTTGAT
Hs.77961	135	GATTACATCGCCCTGAACGAG	136	TCCATCAACCTCTCATAGCAAA
X64707	137	GTAAGATCCGCAGACGTAAGG	138	CTGAAGTCAGCCTCTGTAACTCTC
Hs.6780	139	ACTGAGCCCACTTCTTGTGG	140	ACCTTCTCCCTGTTGCTGTC
Hs.153428	141	TCAAAGTATTTAGCTGACTCGCC	142	TAGTCACTCCAGGTTTATGGAGG
A1066764	143	GGGAACTTGAATTCGTATCCATC	144	CTGAATCTCAAACCTGGAGAGTG
cl.5994	145	GATCATCTTTCCTGTTCCAGAG	146	CTGGAAGGTTCTCAGGCTTTTA
D67025	147	GTACGACCAGGCTGAGAAGC	148	ATCTTCGGGGCTATCCAACCT
x16064	149	TCAGCCACGATGAGATGTTT	150	TGTGGATGACAAGCAGAAGC
M80469	151	ACCTTAGGAGGCGAGTTGGT	152	AGGGGTCAACCTTGAACAG
E02628	153	GCATCCTACCACCACTCGT	154	GCAGCATCACCAGACTTCAA
HUMTHYB4	155	ACAAACCCGATATGGCTGAG	156	GCCAAATGCTTGTGGAATGTA
Hs.116922	157	TCGGACCATAATCCAAGTTACC		
x15940	158	TAACCCGAGAATACACCATCAAC	159	ATGGTTTATTGACGGCAGAAG

[0080]

(Measurement of signal strength of RT-PCR product and calculation of prognosis score)

- 5 The signal strength of the RT-PCR product was measured and evaluated in the same method as described in Example 1, and 10 genes with p values of 0.05 or lower in the t-test were selected as a candidate; of them, expression levels of 3 genes were higher in the 5y-R group than in the 5y-F group. The expression levels of 7 genes were higher in the 5y-F group than in the 5y-R group. Base on this information, the present inventors have tried to
- 10 establish a scoring system for predicting the postoperative prognosis of node-negative breast cancer.

[0081]

For obtaining expression level to be a subject of each gene, the expression ratio (ER) to the GAPDH expression was calculated according to the following formula:

- 15 ER of gene A = 16 bit imaging score of semi-quantitative PCR (strength of band stained with ethidium bromide) of gene A of cancer sample X/16 bit imaging score of GAPDH of gene A of cancer sample X

[0082]

(Definition of scoring system for predicting postoperative prognosis of node-negative breast

cancer)

For obtaining the postoperative gene prognosis index of node-negative breast cancer, prognosis score (PS) was defined; (sum of normalized expression ratios of genes highly expressed in 5Y-R group as compared in 5Y-F group) - (sum of normalized expression ratios of genes highly expressed in 5Y-F group as compared in 5Y-R group)

[0083]

A significance of the expression ratio between two groups was evaluated by the Student's t-test. All statistical methods were carried out by Statview version 5.0 (SAS Institute, Cary, NC).

[0084]

(Result)

Clinicopathological findings of 24 breast cancer patients whose genome-wide gene expressions have been investigated are summarized in Table 3. The present inventors have investigated the gene expression by a cDNA microarray composed of 25344 human genes, for tumors from node-negative breast cancer patients of 12 cases showing survival free of disease for 5 years or more after an operation (5Y-F) and node-negative breast cancer patients of 12 cases showing recurrence of breast cancer within 5 years after a surgical operation (5Y-R). The clinical backgrounds were allowed to coincide in age, tumor diameter, estrogen receptor and progesterone receptor, and pathology between two groups.

[0085]

The data of a cDNA microarray was analyzed by the Mann-Whitney test and the Random-permutation test, and genes showing different expressions between 5Y-R tumor and 5Y-F tumor were identified. Through this filter, 58 genes in total were selected, and of them, 21 genes showed significant strong expression in 5Y-R tumor. 37 genes showed higher expression in 5Y-F tumor.

[0086]

The 37 genes showed higher expression in 5Y-F tumor as compared in 5Y-R tumor had six ESTs and one virtual protein (Table 5A, a difference in expression between groups is

expresses as “foldchange”).

[0087]

(Gene with significant high expression in 5Y-F tumor as compared in 5Y-R tumor)

[0088]

5 [Table 5A]

Ac./HS	kind	fold change	p value
M90439	molecular marker (EPC-1) gene	2.324	0.0014
AF047472	spleen mitotic checkpoint BUB3 (BUB3)	2.889	0.0021
Hs.118251	ESTs	2.121	0.0031
D26125	3 alpha-hydroxysteroid/dihydrodiol dehydrogenase DD4, partial cds	2.084	0.0038
Hs.8619	SRX(sex determining region Y)-box 18	3.375	0.0041
X16135	novel heterogeneous nuclear RNP protein, L protein	4.839	0.0042
Hs.9006	VAMP(vesicle-associated membrane protein)-associated protein A,33kDa	3.807	0.0058
M18963	islet of Langerhans regenerating protein (reg)	2.022	0.0060
X07979	integrin beta 1 subunit	2.997	0.0068
AF018080	PYRIN (MEFV)	4.016	0.0071
Hs.58464	ESTs	5.415	0.0079
S79867	type I keratin 16 [human, epidermal keratinocytes, mRNA Partial, 1422 nt]	2.254	0.0090
J02854	myosin light chain (MLC-2)	2.668	0.0090
Z35309	adenylate cyclase8(brain)	2.264	0.0094
Hs.83097	hypothetical protein FLJ22955	4.979	0.0096
Hs.79137	protein-L isopartate(D-aspartate)-o-methyltransferase	2.401	0.0105
Hs.4864	ESTs	2.043	0.0107
Y00052	Peptidylprolyl isomerase A(cyclophilin A)	2.866	0.0107
Hs.5002	copper chaperone for superoxide dismutase; CCS	2.032	0.0114
U43519	dystrophin-related protein 2 (DRP2)	2.022	0.0114
Hs.106326	ESTs	4.733	0.0123
Hs.94653	neurochondrin(KIAA0607)	2.08	0.0129
M13436	ovarian beta-A-inhibin	2.946	0.0135
X51441	serum amyloid A (SAA) protein partial, clone pAS3-alpha	2.383	0.0155
Hs.108623	thrombospondin 2	2.019	0.0174
Hs.5318	ESTs	4.38	0.0174
Hs.69469	GA17 protein	2.279	0.0197
AA777648	peripheral myelin protein 22	2.386	0.0209
Hs.106131	ESTs	2.022	0.0213
X54079	heat shock protein HSP27	5.637	0.0217
D67025	proteasome (prosome, macropain) 26S subunit, non-ATPase, 3	3.179	0.0359
M80469	MHC class I HLA-J gene	3.572	0.0380
AI041182	ov77e07.x1 Soares_testis_NHT Homo sapiens cDNA clone IMAGE1643364	2.321	0.0380
AA148265	RIBOSOMAL PROTEIN L21.	2.019	0.0440
Hs.4943	Inter-Alpha-Trypsin Inhibitor Heavy Chain LIKE gene	2.426	0.0442
X99920	S100 calcium-binding protein A13	3.326	0.0456
U22970	interferon-inducible peptide (6-16) gene	2.741	0.0465

[0089]

In Table 5B, 21 genes highly expressed in the 5Y-R group are listed. Of them, five
 10 genes are ESTs and one gene encodes a virtual protein. From the panel including 58 genes,
 marker for postoperative prognosis were selected according to the following standard; (1)
 Having higher signal strength than cut off level situated in at least 60% of cases; (2) $|\mu_R - \mu_F|$
 > 1.0 . Here, $\mu_R(\mu_F)$ shows an average value derived from logarithm converted expression
 ratio in the case of 5Y-R or 5Y-F.

15 [0090]

(Gene with significant high expression in 5Y-R tumor as compared in 5Y-F tumor)

[0091]

[Table 5B]

Ac./HS	kind	fold change	p value
X75252	Prostatic Bindig protein	4.506	0.0011
AA989127	major histocompatibility complex, class I,C	5.731	0.0060
Hs.128520	ESTs	1.419	0.0067
HSMLN50	ESTs	3.482	0.0071
AF058701	DNA polymerase zeta catalytic subunit (REV3)	2.185	0.0085
AF043473	delayed-rectifier K ⁺ channel alpha subunit (KCNK1),Potassium voltage-gated channel, delayed-rectifier, subfamily S, member 1	4.786	0.0144
Hs.26052	hypothetical protein MGC43306	4.829	0.0150
Hs.77961	major histocompatibility complex, class I, B	5.775	0.0152
Hs.26484	HLA interacting protein 3	5.07	0.0157
U44798	U1-snrNP binding protein homolog (70kD)	2.615	0.0194
Hs.77961	MHC class I HLA-Bw62	5.775	0.0209
X64707	BBC1 mRNA(ribosomal protein L13)	2.758	0.0210
Hs.6780	PTK9L protein tyrosine kinase 9-like (A6-related protein)	2.749	0.0220
Hs.153428	Es:s	3.164	0.0234
AI066764	lectin, galactoside-binding, soluble, 1. (galectin 1)	2.606	0.0275
c15994	ESTs	2.844	0.0286
x16064	Tumor protein, translationally-controlled 1	3.567	0.0366
E02628	polypeptide chain elongation factor-1 alpha	4.055	0.0427
HUMTHYB4	thymosin beta-4	4.05	0.0436
Hs.116922	ESTs	2.538	0.0494
x15940	ribosomal protein L31.	2.125	0.0499

5

[0092]

7 genes highly expressed in 5Y-F tumor as compared in 5Y-R tumor (Hs.94653, M13436, Hs.5002, D67025, M80469, Hs.4864 and Hs.106326; p = 0.0018, 0.0011, 0.001, 0.008, 0.0081, 0.0018 and 0.001; each according to Student's t-test) and 3 genes relatively
 10 highly expressed in 5Y-R tumor (AF058701, AI066764, and x15940; p = 0.0351, 0.00161 and 0.0001; each according to Student's t-test) coincided with standards, and were selected as a prognosis marker (Table 6).

[0093]

(Genes selected as prognosis marker for node-negative breast cancer)

15 [0094]

[Table 6]

AF058701	DNA polymerase zeta catalytic subunit (REV3)
AI066754	lectin, galactoside-binding, soluble, 1 (galectin 1)
x15940	ribosomal protein L31.
Hs.94653	neurochondrin(KIAA0607)
M13436	ovarian beta-A-inhibin
Hs.5002	copper chaperone for superoxide dismutase; CCS
D67025	proteasome (prosome, macropain) 26S subunit, non-ATPase, 3
M80469	MHC class I HLA-J gene
Hs.4864	ESTs
Hs.106326	ESTs

[0095]

Expressions of these markers were confirmed by a normalized semi-quantitative
5 RT-PCR experiment for GAPDH expression. Fig. 3 shows results of RT-PCR of three
marker genes highly expressed in samples from 12 patients showing recurrence of breast
cancer (5Y-R group). Fig. 4 shows results of 7 marker genes highly expressed in the 5Y-F
group (5 years survival). The expression ratios of these 10 genes were used for definition
of prognosis index.

10 [0096]

Prognosis score (PS) was defined as described below;

PS = (sum of normalized expression ratios of 3 genes highly expressed in 5Y-R tumor)
- (sum of normalized expression ratios of 7 genes highly expressed in 5Y-F tumor)

[0097]

15 The prognosis scores of 24 cases investigated are summarized in Table 7 together with
the expression ratio of each marker gene. The PS system predicted poor prognosis of cases
R1 to R12 having prognosis scores of more than 3. On the other hand, excellent prognosis
was predicted for cases F1 to F12 having scores of lower than -16. The predictions
coincided with actual clinical results of them with an accuracy of 100% (Fig. 5). The
20 average PS of the 5Y-R group was 9.44 and the average PS of the 5Y-F group was -28.92.

[0098]

(Prognosis score for recurrence of node-negative breast cancer)

[0099]

[Table 7]

No.	x1594()	AF058701	AI066764	Hs.5002	Hs.94653	M13436	M80469	D67025	Hs.4864	Hs.106326	PS
1n	8.90	2.70	8.35	1.50	0.82	1.47	2.43	2.72	2.60	2.55	5.86
2n	7.02	2.19	7.48	1.14	0.50	1.51	2.32	1.27	1.89	0.62	7.44
3n	7.57	2.36	10.96	1.40	0.55	2.29	3.51	2.38	1.79	0.44	8.53
4n	8.57	2.79	9.78	1.75	1.42	2.02	3.30	3.03	3.44	3.02	3.16
5n	14.96	2.56	18.01	3.88	0.53	0.67	3.96	2.76	3.78	1.83	18.12
6n	16.94	3.97	12.76	0.11	0.73	1.50	3.19	2.01	3.60	4.41	18.12
7n	14.51	3.02	11.62	0.37	2.24	2.05	2.14	1.45	1.64	2.96	16.30
8n	9.50	2.81	10.43	2.86	1.64	1.95	5.40	3.18	1.89	1.79	4.03
9n	8.29	2.96	8.32	0.78	0.55	1.91	1.50	1.31	1.40	2.80	9.32
10n	6.78	2.06	10.59	0.39	1.93	0.70	2.49	3.56	1.27	0.84	8.25
11n	7.30	1.38	10.89	3.03	2.82	0.46	2.18	3.09	2.00	2.16	3.83
12n	8.60	3.81	15.86	3.31	3.46	0.70	3.19	1.82	2.54	2.95	10.30
1nR	4.67	0.81	4.69	4.13	2.98	3.80	7.78	5.34	7.59	8.47	-29.92
2nR	4.32	0.63	3.88	2.82	2.68	2.89	4.51	3.74	4.86	9.28	-21.95
3nR	10.54	0.56	7.28	2.40	2.06	2.10	8.18	6.02	6.02	8.55	-16.95
4nR	5.59	0.56	4.85	3.22	3.69	2.89	11.18	3.31	6.39	11.36	-31.04
5nR	5.56	0.18	4.97	5.57	4.57	1.15	3.18	4.85	5.56	12.68	-26.85
6nR	4.50	0.51	4.01	6.81	2.54	5.45	6.61	7.49	7.16	6.18	-33.22
7nR	5.09	0.97	4.72	3.14	3.74	5.57	7.95	3.94	7.90	9.71	-31.17
8nR	4.93	0.54	4.46	7.53	4.95	5.93	11.03	1.96	6.21	7.75	-35.43
9nR	5.25	1.17	5.15	3.09	3.33	3.30	10.05	2.66	4.76	10.82	-26.50
10nR	5.36	0.59	5.96	3.67	2.73	2.47	4.66	3.12	10.63	8.27	-23.69
11nR	4.99	1.02	5.71	7.48	4.51	6.22	4.61	4.28	10.65	9.20	-35.23
12nR	4.84	0.30	4.98	7.57	6.07	5.04	7.05	3.07	7.42	8.98	-35.08

[Example 3

5 [0100]

Evaluation of gene expression function for prediction of the postoperative prognosis in primary breast cancer

[0101]

(Tissue sample)

10 A tissue sample was collected in the same manner as described in Example 1. Among 954 patients clinically traced during a period of 5 years or more or until death after an operation for breast cancer in a period from 1995 to 1997, 10 cases of death within 5 years after an operation and 10 cases of survival free of disease for 5 years or more after an operation were selected as a sample. The clinical backgrounds between two patient groups
15 were allowed to coincide as strictly as possible regarding age, metastasis to lymph node, tumor diameter and tissue type (Table 8). The clinical backgrounds of additional 20 cases used for testing the final prediction system are summarized in Table 9.

[0102]

(Clinical profile of patients used for microarray analysis)

20 [0103]

[Table 8]

	Case	T	N	M	Stage	Age	NL ^a	ly ^b	f ^c	ER ^d
<u>Survived</u>	MS1	2	1	0	II	52	4	1	2	P
	MS2	2	2	0	II	47	2	0	1	P
	MS3	2	2	0	II	40	5	0	1	N
	MS4	2	2	0	II	64	3	0	1	N/A
<u>Dead</u>	MD1	2	2	0	II	47	5	0	0	P
	MD2	2	2	0	II	34	3	3	0	N
	MD3	2	2	0	II	66	4	0	3	N
	MD4	2	0	0	II	71	2	0	1	P

a) Number of lymph nodes involved.

b) Lymph vessel invasion: 0, no cancer cells in vessels.
3, many cancer cells in vessels.

c) Fat invasion: 0, no invasion to fat tissue; 3, severe invasion to fat tissue.

d) Estrogen receptor status: P, positive; N, negative; N/A, not available.

[0104]

5 (Clinical profile of patients used for RT-PCR analysis)

[0105]

[Table 9]

	Case	T	N	M	Stage	ly ^a	f ^b
<u>Survived</u>	S1	2	0	0	II	0	1
	S2	2	2	0	II	1	0
	S3	2	2	0	II	0	2
	S4	2	1	0	II	1	0
	S5	2	2	0	II	3	2
	S6	2	0	0	II	0	0
	S7	2	1	0	II	0	0
	S8	2	1	0	II	0	2
	S9	2	1	0	II	1	2
	S10	2	1	0	II	0	0
<u>Dead</u>	D1	2	1	0	II	0	1
	D2	2	2	0	II	0	0
	D3	2	2	0	II	3	0
	D4	2	2	0	II	0	3
	D5	2	2	0	II	1	3
	D6	2	1	0	II	0	1
	D7	2	0	0	II	0	1
	D8	2	1	0	II	0	0
	D9	2	4	0	IV	1	0
	D10	2	1	0	II	0	2

a) Lymph vessel invasion

b) Fat infiltration

	Age ^c	lymph node correlation ^d
<u>Survived</u>	52.8	7.6
<u>Dead</u>	56.0	5.4

c) Mean of age d) Average number of lymph nodes involved

[0106]

(Clinicopathological parameter)

- 5 The clinicopathological parameters were checked by the method described in Example 1.

[0107]

(Preparation of cDNA microarray)

A cDNA microarray was prepared by the method described in Example 2.

[0108]

(RNA extraction and RNA amplification)

5 RNA was extracted using TRIzol (Invitrogen, Carlsbad, CA, USA). For removing degenerate RNA, each extracted RNA (1 μ g) was subjected to electrophoresis on 3.0% formaldehyde denatured gel. For removing DNA mixing, purification was carried out using RNeasy kit (QIAGEN, Valencia, CA). Amplification was carried out based on T7 RNA polymerase base by Messsage Amp aRNA kit (Ambion, Austin, TX), and RNA used
10 for microarray analysis was prepared. In the first amplification, RNA (5 μ g) was used as a template. Thereafter, the firstly amplified RNA (aRNA) (2 μ g) was used as a template for the second amplification. The amplified aRNAs were purified by RNeasy purification kit, and the amount of each aRNA was measured by a spectrophotometer.

[0109]

15 (Labeling of aRNA, hybridization and data analysis)

A hybridization probe was produced using aRNA (5 μ g) for producing fluorescent probe obtained by second amplification, using Amino Allyl-cDNA labeling kit (Ambion, Austin, TX). Probes derived from cancer RNA and normal control RNA were labeled with Cy5 or Cy3 Mono-Reactive Dye (Amersham Bioscience UK Limited, Buckinghamshire,
20 UK), respectively.

[0110]

For removing an unbound dye, a labeled probe was purified by QIA quick PCR purification kit (QIAGEN, Valencia, CA). Each 10 pmol of fluorescent labeled probes from tumor and normal RNA were mixed with 4x microarray hybridization buffer
25 (Amersham (UK)) and de-ionized formamide. The probe mixture was hybridized to a cDNA array at 40°C for 15 hours. Thereafter, the mixture was washed with 0.1x SSC containing 0.2% SDS once for 5 minutes, then, twice for 10 minutes. All procedures were carried out in Automated Slide Processor System (Amersham). The signal strength of each

hybridization was read by Gene Pix 4000 (Amersham), and evaluated by Gene Pix Pro 3.0 (Axon Instruments, Inc., Foster City, CA, USA). The read signals were normalized by the total gene normalization method (Yang, Y.H., Dudoit, S., Luu, P., Lin, D.M., Peng, V., Ngai, J., and Speed, T.P. (2002). *Nucleic Acids Res* 30, e15.; Manos, E.J., and Jones, D.A. (2001). *Cancer Res* 61, 433-438).

[0111]

For confirming genes showing different expressions between a survival group and a dead group, normalized signals were analyzed by the Mann-Whitney test; the normalized signals were applied to a series of Xs. X represents Cy5/Cy3 signal strength ratio for each gene and each sample (Ono, K., et al. (2000). *Cancer Res* 60, 5007-5011). Genes showing a U value of 0 in the Mann-Whitney test and genes showing a difference of 2-fold or more in expression strength between two groups were selected. Genes with S/N ratios of less than 3.0 were excluded from investigation.

[0112]

(Semi-quantitative RT-PCR experiment and gene expression ratio)

For verifying the data of a microarray, the present inventors carried out a semi-quantitative RT-PCR experiment by reverse-transcribing RNA (10 µg). For adjusting the concentration of the transcribed cDNA, GAPDH was selected as an internal control, and semi-quantitative RT-PCR was carried out (Ono, K., et al. (2000). *Cancer Res* 60, 5007-5011). Primers for GAPDH were 5'-ggaaggtgaaggtcggagt-3 (Forward) and 5-tgggtggaatcatattggaa-3 (Reverse). After adjusting the concentration of the primer, semi-quantitative RT-PCR was carried out on selected genes in samples from the survival group and the dead group. Primers for the genes (Table 10) were designed based on sequence information of NCBI Gen Bank (<http://www.ncbi.nlm.nih.gov/>) and primer 3 on website (http://www.genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi). Each semi-quantitative RT-PCR experiment was performed using, as a template, cDNA (1 µl) having been adjusted concentration, 5 U Takara EX Taq (Takara, Otsu, Japan), 1x PCR buffer (10 mM Tris-HCl, 50 mM KCl, 1.5 mM MgCl₂), and 10 nM dNTPs and 10 pmol of forward

and reverse primers, in a total amount of 30 μ l.

SEQ ID No. 160 ggaaggtgaaggtcggagt

SEQ ID No. 161 tgggtggaatcatattggaa

[0113]

5 (Primer of semi-quantitative PCR)

[0114]

[Table 10]

gene	SEQ ID No.	Forward	SEQ ID No.	Reverse
PtIP	162	CCTCCAACTGCTCCTACTCG	163	TCGAAGCCTCTGTGTCCTTT
C1r	164	GAAGTTGTGGAGGGACGTGT	165	GACTTCCAGCAGCTTCCATC
DPYSL3	166	CATGTA CTGAGCAGGCCAGA	167	AAGATCTTGGCAGCGTTTGT
PTK9l.	168	TTGTGATTGAGGACGAGCAG	169	AATGGTTTCCCGCTCTAGGT
CPE	170	CTCCTGAGACCAAGGCTGTC	171	TGAAGGTCTCGGACAAATCC
α -tubulin	172	GGAACGCCTGTCA GTTGATT	173	CTCAAAGCAAGCATTGGTGA
β -tubulin	174	TCTGTTCCGCTCAGGTCCTTT	175	TGGTGTGGTCAGCTTCAGAG
HSP 90-a	176	AAAAATGGCCTGAGTTAAGTGT	177	TCCTCAATTTCCCTGTGTTTG
MDH	178	TGCACACTAACAGCATGACG	179	GAATTTCTTTCCTCTGCCTGA
NDUFE3	180	GGGATAAACCAAGACAAGTAGGC	181	GGACATGAGCATGGACATCA

10 [0115]

For evaluating the strengths of gene expressions between the survival group and the dead group, each semi-quantitative PCR product (8 μ l) was subjected to electrophoresis on 2.5% agarose gel, and stained with ethidium bromide. The concentration of each stained sample was measured by AlphaImager 3300 (Alpha Innotech, San Leandro, CA) using background correction. For obtaining the expression level of each gene, the expression ratio was normalized with the expression level of GAPDH.

[0116]

The expression ratio was defined by the following formula: Expression ratio of gene A = 16 bit imaging score of semi-quantitative PCR (strength of band stained with ethidium bromide) of gene A in cancer sample X/16 bit imaging score of GAPDH in cancer sample X

[0117]

(Definition of prognosis index (PI) of primary breast cancer)

The present inventors defined the prognosis index (PI) of primary breast cancer by subtracting the sum of normalized expression ratios of genes highly expressed in the 5D

group from the sum of normalized expression ratios of genes highly expressed in the 5S group. A significance of expression ratios between two groups was evaluated by the Student's t-test. Comparison of PI between the 5S group and the 5D group was carried out by the Mann-Whitney test. All the statistics were stored using Statview version 5.0
5 (SASInstitute Inc., Cary, NC).

[0118]

(Result)

On a cDNA microarray composed of 18432 human genes, genome-wide gene expression functions of tumors from 8 breast cancer patients were examined. Four patients
10 survived free of disease for 5 year or more after an operation (5S), and four patients died of breast cancer within 5 years after the operation (5D). The clinical backgrounds between two patient groups were allowed to coincide as strictly as possible regarding age, tumor diameter, metastasis to lymph node, hormone receptor condition and tissue type (Table 8).

[0119]

15 For identifying genes showing different expressions between the 5D group and the 5S group, the present inventors analyzed the data of the cDNA microarray by the Mann-Whitney test. 23 genes in total among which six genes are ESTs/virtual proteins are genes showing a U value of 0 in the Mann-Whitney test and highly expressed in the 5S group (Table 11).

20 [0120]

(Gene group highly expressed in survival group by microarray analysis)

[0121]

[Table 11]

Gene name and detail

	Accession Number	Fold change
IMAGE39159 3' similar to gbJ04173 PHOSPHOGLYCERATE MUTASE, BRAIN FORM	R51864	4.304
IMAGE22798 3', mRNA sequence	R39171	2.918
cDNA clone IMAGE:1693352 3', mRNA sequence	A1140851	2.891
CCNDBP1 cyclin D-type binding protein 1	AF082569	3.202
ESTs	A1446435	3.251
pro-alpha-1 type 3 collagen	X14420.1	3.394
complement component C1r	J04080.1	3.396
DPYSL3 dihydropyrimidinase-like 3	D78014	3.625
ribosomal protein L6	X69391.1	3.807
PTK9L protein tyrosine kinase 9-like (A6-related protein)	Y17169.1	4.143
Homo sapiens full length insert cDNA YN88E09	AF075050.1	4.257
somatostatin receptor isoform 2 (SSTR2) gene	M81830.1	5.475
CPE carboxypeptidase E	NM_001873.1	5.807
YR-29 hypothetical protein YR-29	AJ012409.1	6.333
IMAGE4822062, mRNA	BC034811	6.373
KIAA1832 protein, partial cds	AB058755.1	13.352
CREG cellular repressor of E1A-stimulated genes	AF084523.1	2.739
Homo sapiens putative splice factor transformer2-beta mRNA, complete cds	U61267.1	2.55
Human N-acetyl-beta-glucosaminidase (HEXB) mRNA, 3' end	M13519.1	2.698
Human cytochrome b5 mRNA, complete cds	M22865.1	2.881
Human p52 mRNA induced by estrogen from human breast cancer cell line MCF-7	X00474.1	2.702
Human alpha-tubulin mRNA, complete cds	K00558	4.655
Homo sapiens clone 24703 beta-tubulin mRNA, complete cds	AF070561.1	3.917

[0122]

Table 12 describes 21 genes highly expressed in general in the 5D tumor, including 6
5 ESTs/virtual proteins, and having a U value of 0 in the Mann-Whitney test. In the table, a
difference in gene expression between two groups is shown as “foldchange”.

[0123]

(Gene group highly expressed in dead group by microarray analysis)

[0124]

10 [Table 12]

Gene name and detail	Accession Number	Fold change
Lyam-1 mRNA for leukocyte adhesion molecule-1	X16150.1	7.459
APM2 adipose specific 2	NM_006829.1	4.853
DNA polymerase gamma mRNA, nuclear gene encoding mitochondrial protein	U60325.1	4.269
FLJ22128 fis, clone HEP19543	AK025781	4.109
actin related protein 2/3 complex, subunit 4, 20kDa (ARPC4)	NM_005718.2	4.058
Scd mRNA for stearyl-CoA desaturase	AB032261.1	3.794
novel heterogeneous nuclear RNP protein, L protein	X16135.1	3.771
ENSA endosulfine alpha	AF157509.1	3.511
IMAGE:26483 5' similar to gb:X15183_cds1 HEAT SHOCK PROTEIN HSP 90-ALPHA	R12732	3.086
malonyl-CoA decarboxylase (MLYCD)	NM_012213	3.067
anion exchanger 3 brain isoform (bAE3)	U05596.1	2.889
IMAGE:43550 3', mRNA sequence	H05914	2.345
cDNA FLJ23636 fis, clone CAS07176.	AK074216	2.426
IMAGE:26366 3' similar to gb:D16234 PROBABLE PROTEIN DISULFIDE ISOMERASE ER-60 PRECURSOR	R20554	2.519
Similar to hypothetical protein PRO2831, clone MGC:23813 IMAGE:4273837, mRNA, complete cds	BC017905.1	2.551
FLJ40629 hypothetical protein FLJ40629	AK097948.1	2.417
ribosomal protein L29 (humrpl29) mRNA, complete cds	U10248.1	2.203
EST, clone IMAGE:745452, 3'end	AA625869	2.591
KIAA1554 KIAA1554 protein	AB046774.1	2.544
IMAGE:53316 3' similar to SP:MDHC_MOUSE P14152 MALATE DEHYDROGENASE, CYTOPLASMIC	R15814	2.867
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 3 (12kD, B12), clone MGC:9039 IMAGE:3881592	BC018183	4.972

[0125]

From 23 genes highly expressed in the 5S group and 21 gene highly expressed in the 5D group, prediction markers for postoperative prognosis were selected according to the following standards; (1) In microarray analysis, a difference in the signal strength between 5S and 5D is larger than 2.0-fold in all cases; (2) The signal strength differs significantly between 5S and 5D in semi-quantitative PCR (p value < 0.05 in Student's t-test); (3) The result of semi-quantitative PCR was re-confirmed by independent triple experiments. 7 genes highly expressed in the 5S tumor and 3 genes highly expressed in the 5D tumor satisfied these standards for selecting a prognosis marker.

[0126]

7 genes highly expressed in the 5S group are constituted of genes encoding pro-alpha-1 type 3 collagen (PIIIP), complement component C1r, dihydropyrimidinase-like

3 (DPYSL3), proteintyrosinekinase 9-like (PTK9L), carboxy peptidase E (CPE), α -tubulin and β -tubulin. The p values in the Student's t-test of these marker genes were 0.00039, 0.0012, 0.0042, 0.036, 0.039, 0.034 and 0.00069, respectively.

[0127]

5 3 marker genes highly expressed in the 5D group encoded heat shock protein HSP 90-alpha gene, malatedehydrogenase, and NADH dehydrogenase (ubiquinone) 1 beta subcomplex 3 (NDUFB3). The p values in the Student's t-test of these genes were 0.05, 0.0055 and 0.011, respectively.

[0128]

10 The present inventors normalized the experiment results of semi-quantitative RT-PCR by GAPDH as an internal control and evaluated the results, verifying selection of marker genes.

[0129]

15 The present inventors carried out semi-quantitative PCR for checking additional 20 cases randomly selected. 10 of these patients died of breast cancer within 5 years after an operation, and remaining 10 patients survived free of disease for 5 years or more after the operation. Fig. 7 shows the results of RT-PCR of 7 marker genes highly expressed in the 5S tumor. Fig. 8 shows the results of RT-PCR of 3 marker genes highly expressed in the 5D tumor.

20 [0130]

 The present inventors defined the prognosis index (PI) as described below: (sum of normalized expression ratios of genes highly expressed in 5S group) - (sum of normalized expression ratios of genes highly expressed in 5D group). The expression ratios of the selected marker genes are summarized together with prognosis indices for further test
25 examples in Table 13.

[0131]

(Expression ratio of gene and prognosis index)

[0132]

[Table 13]

	Gene highly expressed in 5S					Gene highly expressed in 5D							
	PIIP	C1r	DPYSL3	PTK9L	CPE	A-tubulin	B-tubulin	HSP 90	MDH	NDUFB3	Sum of S *	Sum of D **	PI +
S1	1.3	4.0	2.1	3.3	2.4	0.8	2.5	1.5	0.2	1.5	16.9	3.2	13.7
S2	5.7	3.5	3.3	3.4	6.0	1.2	5.1	2.2	0.6	2.2	28.1	5.0	23.1
S3	3.1	5.8	2.2	3.4	8.1	1.8	5.3	1.5	0.4	1.5	29.7	3.5	26.2
S4	7.1	10.2	8.6	6.0	16.0	4.1	8.0	3.4	4.8	3.4	60.0	11.5	48.5
S5	6.8	7.4	7.2	6.9	11.2	2.7	7.0	5.5	3.6	5.5	49.1	14.6	34.5
S6	4.0	4.2	1.7	2.2	3.6	0.9	6.0	2.9	0.9	2.9	22.7	6.6	16.1
S7	2.3	4.0	1.1	1.6	0.6	0.7	3.4	0.4	0.3	0.4	13.7	1.1	12.6
S8	3.3	3.6	1.1	0.7	0.8	1.3	5.0	2.3	1.4	2.3	15.9	6.0	9.8
S9	3.1	3.9	2.7	3.7	2.9	1.6	4.1	1.0	1.2	1.0	21.9	3.2	18.8
S10	2.9	3.0	0.9	1.5	1.2	1.0	1.7	1.3	0.4	1.3	12.2	3.0	9.2
D1	0.1	2.9	0.4	1.9	2.9	0.7	0.8	3.4	3.0	3.4	9.6	9.7	-0.1
D2	0.2	0.6	0.1	0.2	0.8	0.2	0.8	1.0	4.9	1.0	2.9	7.0	-4.1
D3	0.2	3.7	0.2	1.0	0.6	0.6	2.8	3.6	6.6	3.6	9.0	13.8	-4.8
D4	0.2	1.4	0.4	0.9	1.0	0.5	1.7	3.5	3.6	3.5	6.1	10.7	-4.6
D5	0.1	1.3	0.1	0.9	0.6	0.5	1.0	3.2	0.3	3.2	4.5	6.7	-2.2
D6	2.2	2.5	1.2	1.9	2.0	0.5	1.7	3.8	3.5	4.2	12.0	11.5	0.5
D7	2.2	2.1	0.9	1.9	2.4	0.3	1.6	1.9	1.4	2.0	11.5	5.3	6.2
D8	1.6	2.7	1.1	2.6	1.8	0.4	1.8	3.4	2.8	3.4	12.0	9.6	2.5
D9	1.2	1.4	0.6	1.6	1.2	0.6	2.4	2.2	0.7	2.2	9.2	5.0	4.1
D10	0.5	0.8	0.4	0.6	0.4	0.4	1.3	3.6	1.6	3.6	4.5	8.9	-4.4

* Sum of ER of PIIP, C1r, DPYSL3, CPE, α and β -tubulin

** Sum of ER of HSP 90, MDH and NDUFB3

+PI: Sum of S - sum of D

[0133]

5 PI predicted correctly the actual clinical results of higher prognosis indices (> 7) of 10 cases (S1 to S10) in total in the 5S group and prognosis indices (< 7) of 10 cases (D1 to D10) in total in the 5D group. PI of the 5S group was 21.2. PI of the 5D group was -0.7. Here, by a PI value of 7 the 5S tumor and the 5D tumor were apparently distinguished ($p = 0.0002$).

10 [INDUSTRIAL APPLICABILITY]

[0134]

The postoperative prognosis prediction system of the present invention is effective for prediction of postoperative risk of a breast cancer patient. Further, the wide-range gene expression list of breast cancer correlated genes of the present invention can provide various

information on progress of breast cancer, and a latent target molecule for breast cancer therapy was predicted by the list.

[BRIEF DESCRIPTION OF DRAWINGS]

[0135]

5 [Fig. 1] Fig. 1 shows a gene group (A) manifesting increase and a gene group (B) manifesting decrease in expression in 5y-D group as compared with 5y-S group.

[Fig. 2] Fig. 2 shows analysis results of semi-quantitative RT-PCR of RNAs derived from 5y-S group and 5y-D group.

[Fig. 3] Fig. 3 shows prognosis scores in individual patients.

10 [Fig. 4] Fig. 4 shows analysis results of semi-quantitative RT-PCR of RNAs derived from 5Y-F group and 5Y-R group.

[Fig. 5] Fig. 5 shows analysis results of semi-quantitative RT-PCR of RNAs derived from 5Y-F group and 5Y-R group.

[Fig. 6] Fig. 6 shows prognosis scores in individual patients.

15 [Fig. 7] Fig. 7 shows analysis results of semi-quantitative PCR of 7 genes highly expressed in 5S tumor.

M: marker ladder

S1-S10: newly inspected tissues of patients survived free of disease for 5 years or more after operation.

20 D1-D10: newly inspected cases of patients died of breast cancer within 5 years after operation.

Difference in expression strength was evaluated by Student's t-test; when p value is 0.05 or less, statistical significance is believed to be present.

[Fig. 8] Fig. 8 shows analysis results of semi-quantitative PCR of 3 genes highly
25 expressed in 5D group. For explanation of marks, please see explanation in Fig. 7.

[Fig. 9] Fig. 9 shows results illustrating prognosis indices (PI) of newly inspected 20 cases. The indices of all 10 patients survived free of disease for 5 years or more were higher than 7. On the other hand, the indices of patients died of breast cancer within 5


years after operation were lower than 7. Distribution of two groups is statistically significant ($p = 0.0002$).

[Sequence Listing]


[Title of Document] ABSTRACT

[Problem] It is intended to provide a system of predicting the postoperative prognosis in a patient with breast cancer from the viewpoint of gene expression based on the data obtained by genome-wide and comprehensive analysis on gene expression in breast cancer.


- 5 [Solution] Expression of human genes is comprehensively analyzed by using a DNA microarray and gene expression functions in various breast cancer conditions are compared, thereby establishing a system of predicting the postoperative prognosis of breast cancer.




[Fig. 2]




[Fig. 3]




[Fig. 4]




[Fig. 5]



[Fig. 6]



[Fig. 7]



[Fig. 8]

p VALUE

M S1 S2 S3 S4 S5 S6 S7 S8 S9 S10 D1 D2 D3 D4 D5 D6 D7 D8 D9 D10

HSP 90- α

0.05


MDH

0.0055


NDUFB3

0.011

GAPDH



[Fig. 9]



SEQUENCE LISTING

<110> Nippon Medical School,
Mitsubishi Kagaku Bio-Clinical Laboratories, Inc.,
Mitsubishi Rayon Co., Ltd.

<120> Genes involved in predicting postoperative prognosis breast cancers

<130> NP04-1001

<160> 181

<170> PatentIn version 3.1

<210> 1

<211> 19

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 1

ggaaggtgaa ggtcggagt

19

<210> 2

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 2

tgggtggaat catattgga

20

<210> 3

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 3

acacttcac tgctccctca tag

23

<210> 4

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 4

ctgcctagac ctgaggactg tag

23

<210> 5

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 5

actgaggcct ttggtagtc g

21

<210> 6
<211> 24
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 6
tctctttatt gtgatgctca gtgg

24

<210> 7
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 7
aaatccttct cgtgtgttga ctg

23

<210> 8
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 8
cagtcattgag ggctaaaaac tga

23

<210> 9

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 9

gaagacaaca agttttaccg gg

22

<210> 10

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 10

atggttttat tgacggcaga ag

22

<210> 11

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 11

aggacacgtc ctctcctctc tc

22

<210> 12
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 12
taaagctagc gaaggaacgt aca

23

<210> 13
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 13
tcccttctgt ttcctcagtg tt

22

<210> 14
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 14
cctgccccga taaaaatatc tac

23

<210> 15
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 15
ttgaccttaa gccctcttttc ctc

23

<210> 16
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 16
ataacgtaca ttcccatgac acc

23

<210> 17
<211> 21
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 17
actttcaaga tgggaccaag g

21

<210> 18

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 18

atatacacag aagcatgacg cag

23

<210> 19

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 19

ttgctggact ctgaaatattc cc

22

<210> 20

<211> 24

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 20

ttcccctgta cagtatttca ctca

24

<210> 21

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 21

ctgagcaatc tgcctatcc tct

23

<210> 22

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 22

gttcagatt cgtgagaatg act

23

<210> 23

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 23

accagtaaca actgtgggat gg

22

<210> 24

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 24

caaatgagct acaacacaca agg

23

<210> 25

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 25

ccccctccac cttgtacata at

22

<210> 26

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 26

gttttcgttt ggcctggtgt g

21

<210> 27

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 27

gtctgagatt ttactgcacc g

21

<210> 28

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 28

attgctaagg ataagtgtg ctc

23

<210> 29

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 29

tgtcagtata gaagcctgtg ggt

23

<210> 30

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 30

ttcttaggcc atcccttttc tac

23

<210> 31

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 31

gcatctgaat gtctttctcc cta

23

<210> 32

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 32

ccataggatc ttgactccaa cag

23

<210> 33

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 33

actgggagtg gaggaatta gag

23

<210> 34

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 34

ctaagtgaag ctccattggg atg

23

<210> 35

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 35

caaactgcaa aatagctccc taa

23

<210> 36

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 36

aggtaaccca aagtgacaaa cct

23

<210> 37

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 37

aagactaaga gggaaaatgt ggg

23

<210> 38

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 38

aggtaaccca aagtgacaaa cct

23

<210> 39

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 39

ttaagtgagt ctccctggct gag

23

<210> 40

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 40

agggcccccta tatccaatac cta

23

<210> 41

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 41

agtcattcag aagccattga gac

23

<210> 42

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 42

tgggtggaat catattggaa

20

<210> 43

<211> 18

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 43

gaaaggtgaa ggtcggagt

18

<210> 44

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 44

tgggtggaat catattggaa

21

<210> 45

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 45

ccagacatcc atggtaccta taa

23

<210> 46

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 46

tatgcattga aaccttacag ggg

23

<210> 47

<211> 24

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 47

ctgttaaca aagcgaggtt aagg

24

<210> 48

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 48

gggttcgca tctcgtttat tag

23

<210> 49

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 49

gacacatagc tcataggcac aca

23

<210> 50

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 50

ttctgggtaca tggtaagtc tca

23

<210> 51

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 51

tccgccatat tgattctgct ta

22

<210> 52

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 52

gtttgctttc tggaccatgg ata

23

<210> 53

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 53

gataacaact ggaccacatc cc

22

<210> 54

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 54

aacaggcaga cgaggtagac ac

22

<210> 55

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 55

gagaaggatg ggtccaccag t

21

<210> 56

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 56

gtacatgggc agcacaaatg tat

23

<210> 57

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 57

atttcattgg tagtatggcc cac

23

<210> 58

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 58

ataccatggg acaggattgt aag

23

<210> 59

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 59

gctcagacca gctcatactt cat

23

<210> 60
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 60
ccaaagactg ggtaggtaa aac

23

<210> 61
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 61
ctggcgcttt ctatcacctc ttc

23

<210> 62
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 62
gactagtgtg aaacaagatg ggc

23

<210> 63
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 63
cttgaaccca ggagtttgag ac

22

<210> 64
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 64
gtgcctcagc ttctgagta gc

22

<210> 65
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 65
ctgggtgctga ctatccagtt ga

22

<210> 66
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 66
ctggtaaact gtccaaaaca agg

23

<210> 67
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 67
ctcttacctg gacaagggtgc gt

22

<210> 68
<211> 21
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 68
ggatgagctc tgc:ccttga g

21

<210> 69

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 69

caatgtttga ccagtcccag a

21

<210> 70

<211> 24

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 70

catgttgtct cagtcctcta ttgg

24

<210> 71

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 71

ggacagcagc tggagtacac a

21

<210> 72
<211> 21
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 72
aatcagattt gtcggtgcct t

21

<210> 73
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 73
ggctctgcac taagaacaca gag

23

<210> 74
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 74
acaactagct ctcagttcag gca

23

<210> 75

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 75

tggagcagta tgacaagcta caa

23

<210> 76

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 76

aagcagcact gcataaactg ttc

23

<210> 77

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 77

taagtacttt cctgtgggtc gct

23

<210> 78
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 78
ccacaaacag gaagctatgt tct

23

<210> 79
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 79
gtactattag ccatggtcaa ccc

23

<210> 80
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 80
ctacagaagg aatgatctgg tgg

23

<210> 81
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 81
atcagtacgg ggaccittaca aac

23

<210> 82
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 82
cctgtactga gctctccaaa gac

23

<210> 83
<211> 20
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 83
tccctagctt cctctccaca

20

<210> 84

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 84

agaatcatgc ctccccttct

20

<210> 85

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 85

accctcaag tgaaggaac tg

22

<210> 86

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 86

ggatcaagag tglgtgtgtg tgt

23

<210> 87
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 87
caatgccaga gagaatatcc aga

23

<210> 88
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 88
gataccatt gtgtaccctc tcc

23

<210> 89
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 89
ccactccaca taaggggttt ag

22

<210> 90
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 90
gaggttctag ctaagtgcag ggt

23

<210> 91
<211> 25
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 91
ccattgacat tggagttaag tatgc

25

<210> 92
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 92
ggcaaagacc acaatttagca at

22

<210> 93
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 93
gaaagcctat gtgaaaagct ggt

23

<210> 94
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 94
ttgtttccag gcattaagtg tg

22

<210> 95
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 95
gcatcttagt ccacacagtt ggt

23

<210> 96

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 96

gcccttacag gtggagtatc ttc

23

<210> 97

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 97

ctcatagcca gcatgacttc ttt

23

<210> 98

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 98

ggttcacttg tgactggcca tct

23

<210> 99
<211> 22
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 99
acttttc tga gcagacgtcc ag

22

<210> 100
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 100
tatcaaaaga acacacaggt ggc

23

<210> 101
<211> 23
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 101
acgttat tcc cagttcctaa acc

23

<210> 102

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 102

agtcctcgggt gactcaatat gaa

23

<210> 103

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 103

agttgaaccc aggtaccttt ctc

23

<210> 104

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 104

ctaggccctt ttagaaaaca tgg

23

<210> 105

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 105

tactgggaac gactaaggac tca

23

<210> 106

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 106

tgctgtgttg agtaggttc tga

23

<210> 107

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 107

tgagagtcct cagagggtat cag

23

<210> 108

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 108

cttgaagtca agagtcctgg tgt

23

<210> 109

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 109

tttctgttg caagttgctg

20

<210> 110

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 110

ccctttaagc ccacttcctc

20

<210> 111

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 111

gatgagaaga tgaagagctt gga

23

<210> 112

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 112

gaggaagctt tatttgggaa gag

23

<210> 113

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 113

acttccctct ctgcctttct g

21

<210> 114

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 114

cagattgttt tggcttctc act

23

<210> 115

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 115

gtctggtcag ctttgcttcc

20

<210> 116

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 116

ggcaagttct gcacagatga

20

<210> 117

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 117

cagctcagtg caccatgaat

20

<210> 118

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 118

gtgggactga gatgcaggat

20

<210> 119

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 119

cacggactca tgaatgtagt gaa

23

<210> 120

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 120

gtgtagtggc acgatcatag ctt

23

<210> 121

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 121

gggaccaaac agaccaaaga

20

<210> 122

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 122

cacccacag agcctgtatt

20

<210> 123

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 123

cggaaggca ctatttcaca at

22

<210> 124

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 124

acaggccac aggtttgtaa c

21

<210> 125

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 125

aagctcttca gctgcgtctc

20

<210> 126

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 126

cctcctccctt ttcagctgtg

20

<210> 127

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 127

tctggaaccc taaaagtgtc gt

22

<210> 128

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 128

tctttcaaca tctctccacc cta

23

<210> 129

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 129

agatacctgg agaacgggaa g

21

<210> 130

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 130

ggaagtaaga agttgcagct cag

23

<210> 131

<211> 18

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 131

attaggtttc acccaaag

18

<210> 132

<211> 19

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 132

agacgagact tgttttctc

19

<210> 133

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 133

cagggacttg gtcacagggtt

20

<210> 134

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 134

ttcttctccc tccccttgat

20

<210> 135

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 135

gattacatcg ccctgaacga g

21

<210> 136

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 136

tccatcaacc tctcatagca aa

22

<210> 137

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 137

gtaagatccg cagacgtaag g

21

<210> 138

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 138

ctgaagtcag cctctgtaac ctc

23

<210> 139

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 139

actgacccca cttcttgtgg

20

<210> 140

<211> 19

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 140

acccttcct gttgctgc

19

<210> 141

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 141

tcaaagtatt tagctgactc gcc

23

<210> 142

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 142

tagtcactcc aggtttatgg agg

23

<210> 143

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 143

gggaacttga attcgatcc atc

23

<210> 144

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 144

ctgaatctca aacctggaga gtg

23

<210> 145

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 145

gatcatcttt cctgttccag ag

22

<210> 146

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 146

ctggaagggt ctcaggtctt ta

22

<210> 147

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 147

gtacgaccag gctgagaagc

20

<210> 148

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 148

atcttcgggg ctatccaact

20

<210> 149

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 149

tcagccacga agagatgttc

20

<210> 150

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 150

tgtggatgac aagcagaagc

20

<210> 151

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 151

accttaggag ggcagttggt

20

<210> 152

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 152

aggggtcaca cttgaacag

20

<210> 153

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 153

gcatcctacc accaactcgt

20

<210> 154

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 154

gcagcatcac cagacttcaa

20

<210> 155

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 155

acaaacccga tatggctgag

20

<210> 156

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 156

gccaatgctt gtggaatgta

20

<210> 157

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 157

tcggaccata atccaagtta cc

22

<210> 158

<211> 23

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 158

taacccgaga atacaccatc aac

23

<210> 159

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 159

atggttttat tgacggcaga ag

22

<210> 160

<211> 19

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 160

ggaaggtgaa ggicggagt

19

<210> 161

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 161

tgggtggaat catattgga

20

<210> 162

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 162

cctccaactg ctctactcg

20

<210> 163

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 163

tcgaagcctc tglgtccttt

20

<210> 164

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 164

gaagttgtgg agggacgtgt

20

<210> 165

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 165

gacttcagc agcttccatc

20

<210> 166

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 166

catgtactga gcaggccaga

20

<210> 167

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 167

aagatcttgg cagcgtttgt

20

<210> 168

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 168

ttgtgattga ggacgagcag

20

<210> 169

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 169

aatggtttcc cgtcttaggt

20

<210> 170

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 170

ctcctgagac caaggctgtc

20

<210> 171
<211> 20
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 171
tgaaggtctc ggacaaatcc

20

<210> 172
<211> 20
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 172
ggaacgcctg tcagttgatt

20

<210> 173
<211> 20
<212> DNA
<213> Artificial

<220>
<223> synthesized

<400> 173
ctcaaagcaa gcattggatga

20

<210> 174

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 174

tctgttcgct caggtccttt

20

<210> 175

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 175

tgggtgtggc agcttcagag

20

<210> 176

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 176

aaaaatggcc tgagttaagt gt

22

<210> 177

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 177

tcctcaattt ccctgtgtt g

21

<210> 178

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 178

tgcacactaa cagcatgacg

20

<210> 179

<211> 21

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 179

gaatttcctt cctctgcctg a

21

<210> 180

<211> 22

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 180

gggataaacc agacaagtag gc

22

<210> 181

<211> 20

<212> DNA

<213> Artificial

<220>

<223> synthesized

<400> 181

ggacatgagc atggacatca

20